

Scotts
spot
dandelion
control

DIRECTIONS

Apply Scotts Spot Dandelion Control to lawn weeds in lawns. For lawn care information contact Scotts Chemical Corporation, P.O. Box 187, Marysville, Ohio 43040.

WALK WEEDS: BEFORE AND DURING THE first mowing, apply above ground liquid to the top of the growing weed. The liquid penetrates the leaf and is carried down the stems to the roots.

- Apply Scotts Spot Dandelion Control to walk weeds before the lawn is mowed.

- Apply Scotts Spot Dandelion Control to lawn weeds before the lawn is mowed.

- For treatment of lawn weeds, apply Scotts Spot Dandelion Control to the weeds before the lawn is mowed. Apply the liquid to the top of the growing weed. The liquid penetrates the leaf and is carried down the stems to the roots.

Apply Scotts Spot Dandelion Control to lawn weeds before the lawn is mowed. Apply the liquid to the top of the growing weed. The liquid penetrates the leaf and is carried down the stems to the roots.

Scotts
spot
dandelion
control

controls broadleaf and
vining weeds in lawns



ACTIVE INGREDIENTS:
Alkanol amine salts (ethanol and isopropanol series) of 2,4-Dichlorophenoxyacetic acid* 1.86%
Dicamba — dimethylamine salt of (2-Methoxy-3,6-dichlorobenzoic acid)** 0.31%
INERT INGREDIENTS 98.03%
Total 100.00%

*equivalent to 1.00% 2,4-Dichlorophenoxyacetic acid.
**equivalent to 0.25% 2-Methoxy-3,6-dichlorobenzoic acid.
EPA Reg. No. 538-48
EPA Est. 5690-IL-1

Distributed by O. M. Scott & Sons
Marysville, Ohio 43040

CAUTION: KEEP OUT OF
REACH OF CHILDREN.
See other side for additional cautions

NET WT.
1 LB. 4 OZ.

Controls all these lawn weeds

- Buckhorn
- Clover
- Crabgrass
- Dandelion
- Field Bindweed
- Henbit
- Plantain
- Redtop
- Sheep Sorrel
- Spotted Spurge
- St. John's Wort
- Syntherisma
- Thick-knee
- Trifolium
- Wedge
- White Clover
- Yarrow

Apply Scotts Spot Dandelion Control for nonselective control of undesirable weeds and grasses.

Scotts promise to you satisfaction or your money back.

MAR 31 1975

Scott's
spot
dandelion
control

DIRECTIONS

For easy control of broadleaf and vining weeds in lawns. Apply anytime weeds are actively growing.

SHAKE WELL BEFORE AND DURING USE. Hold container 2-3 feet above ground. Invert can and direct spray at center of weed. Press actuator firmly—only a light burst needed! Drenching does not increase effectiveness.

- Weeds start to wilt in a few days, gradually decline. A full month may be required for complete fadeout. Repeat in three or four weeks if weeds persist.

- It is not necessary (or desirable) to remove dead weeds. They will eventually disintegrate and disappear if left alone.

- For treatment of entire lawn, use Scott's KANSSEL TURF BUILDER PLUS 2 or BONUS TYPE B (except St. Augustine or dichondra lawns).

CAUTION: Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water. For eyes get medical attention. Wash after each use.


Avoid drift or application to nearby garden plants, flowers or shrubs. Do not apply within dropline of desirable plants. Do not contaminate water used for irrigation or domestic purposes. Do not store near fertilizer, seeds or other pesticides.

Contents under pressure. Do not use near fire or open flame. Store in cool place (not over 120 F) but protect against freezing. Never throw in fire or puncture this container.

Product of U.S.A. O.M. Scott & Sons, Maysville, Ohio 43040

Scott's
spot
dandelion
control

controls broadleaf and
vining weeds in lawns



ACTIVE INGREDIENTS
Allyl amine salts (technical and isopropyl salts) of 2,4-Dichlorophenoxyacetic acid 7.56%
Dichlorophenoxyacetic acid (2-Methoxy-3,6-Achlorobenzoyl acid) 0.31%
Inert ingredients 92.13%
Total 100.00%

Equivalent to 1.00% 2,4-Dichlorophenoxyacetic acid
Equivalent to 0.25% 2-Methoxy 3,6-dichlorophenoxyacetic acid
EPA Reg. No. 538-48
EPA Est. 5590-1E-1
Distributed by O.M. Scott & Sons, Maysville, Ohio 43040

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other side for additional cautions

NET WT. 1 LB. 4 OZ.

Controls all these lawn weeds

- ~~black medic~~
- ~~buckhorn~~
- ~~Canada thistle~~
- ~~chickweed~~
- ~~clover~~
- ~~cudweed~~
- ~~curly dock~~
- ~~dandelion~~
- ~~dollar weed~~
- ~~English daisy~~
- ~~floxer~~
- ~~heavenly~~
- ~~Florida pusley~~
- ~~ground ivy~~
- ~~heavenly~~
- ~~henbit~~
- ~~knockweed~~
- ~~lamb's quarters~~
- ~~marest weed~~
- ~~mustard~~
- ~~penstemon~~
- ~~peppercorn~~
- ~~pigweed~~
- ~~plantain~~
- ~~poison ivy~~
- ~~prickly lettuce~~
- ~~purple~~
- ~~sheep sorrel~~
- ~~shepherd's purse~~
- ~~yellow~~

Also use Scott's Spot Grass and Weed Control for nonselective control of undesirable weeds and grasses.

Scott's promise to you—satisfaction or your money back.

©1975 O.M. Scott & Sons "GOAR"

3 993

MAR 31 1975

534 78

Scott's
**spot
 dandelion
 control**

DIRECTIONS

Apply Scott's Spot Dandelion Control to dandelions in lawns. Apply to the soil around the base of the weed. Do not apply to grass. Do not apply to other plants. Do not apply to lawns that have been recently fertilized. Do not apply to lawns that have been recently watered. Do not apply to lawns that have been recently mowed. Do not apply to lawns that have been recently raked. Do not apply to lawns that have been recently aerated. Do not apply to lawns that have been recently dethatched. Do not apply to lawns that have been recently overseeded. Do not apply to lawns that have been recently sodded. Do not apply to lawns that have been recently topdressed. Do not apply to lawns that have been recently scarified. Do not apply to lawns that have been recently verticillated. Do not apply to lawns that have been recently rolled. Do not apply to lawns that have been recently watered. Do not apply to lawns that have been recently mowed. Do not apply to lawns that have been recently raked. Do not apply to lawns that have been recently aerated. Do not apply to lawns that have been recently dethatched. Do not apply to lawns that have been recently overseeded. Do not apply to lawns that have been recently sodded. Do not apply to lawns that have been recently topdressed. Do not apply to lawns that have been recently scarified. Do not apply to lawns that have been recently verticillated. Do not apply to lawns that have been recently rolled.

Scott's
**spot
 dandelion
 control**

controls broadleaf and
 vining weeds in lawns


ACTIVE INGREDIENTS:
 Alkylamine salts (amine and isopropyl ester) of 2,4-Dichlorophenoxyacetic acid... 1.00%
 Mixture — (ammonium salt of 2-Methoxy-3,6-Dichlorophenoxyacetic acid)*... 0.31%
NET INGREDIENTS: 00.69%
 Total 100.00%

*equivalent to 1.00% 2,4-Dichlorophenoxyacetic acid.
 **equivalent to 0.25% 2-Methoxy-3,6-Dichlorophenoxyacetic acid.
 EPA Reg. No. 836-48
 EPA Est. 4500-B-1
 Distributed by P. M. Scott & Son
 Marysville, Ohio 43040

CAUTION: KEEP OUT OF REACH OF CHILDREN
 See other side for additional cautions

NET WT. 1 LB. 4 OZ.

Controls all these lawn weeds



Controls all these lawn weeds: dandelion, clover, chickweed, crabgrass, goosegrass, knotweed, lambsquarters, pigweed, redtop, smartweed, spurge, and many others.