

wee control  
for grass lawns

**O-X-D**

Treats 5,000 sq ft / \$4.95

6040

**O-X-D**

**weed control  
for grass lawns**

**weeds controlled**

|                |                      |              |
|----------------|----------------------|--------------|
| black medic    | ground ivy           | oxalis       |
| bur clover     | henbit               | pennywort    |
| chickweed      | honeysuckle          | purslane     |
| daisy fleabane | knotweed (sprouting) | white clover |

**When to use**

Apply when weeds to be controlled are growing actively.  
 Apply to well-moistened foliage, ideally in morning or evening  
 when grass is wet with dew.  
 When temperatures are regularly 90° or above, do not apply to  
 predominately bentgrass or bermuda lawns.  
 On newly seeded, sodded or sprigged lawns wait until new  
 grass has been mowed 3 or 4 times.  
 This product will NOT control weedy grasses such as crabgrass.  
 Nor does it have any effect on rosette-type weeds such as  
 dandelions.

Full directions for  
Guide to Successful Use


Use Scotts Spreader (Standard or Automatic) setting #4 1/2  
to ensure proper distribution

On bentgrass or bermuda lawns, make two  
applications three weeks apart using setting #3 1/2

*Feb. 13, 1973*

*538-54*

**O-X-D** weed control




# O-X-D


**weed control  
for grass lawns**

**controls oxalis and other vining  
weeds (see list on other side)**

**DO NOT USE ON DICHONDRA OR ST AUGUSTINE LAWNS**

A PROBLEM SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE





# O-X-D


**weed control  
for grass lawns**

**controls oxalis and other vining  
weeds (see list on other side)**

NOT USE ON DICHONDRA OR ST AUGUSTINE LAWNS

A PROBLEM-SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE.

W



7088

7088

7088

NET WEIGHT 25 1/2 LBS

7088

7088

10 1/2 x 1 1/2

ProTurf

monocot weed control

8802

S

*Scotts*

# ProTurf

## monocot weed control

selective control for monocot weeds listed on back

**KEEP OUT OF REACH OF CHILDREN**  
 Handle with care. Do not inhale dust. Do not get in eyes. If in eyes, flush with water. If on skin, wash with soap and water. Do not eat, drink, or smoke while using. Do not use near water. Do not use on lawns. Do not use on golf courses. Do not use on tennis courts. Do not use on baseball diamonds. Do not use on soccer fields. Do not use on football fields. Do not use on baseball fields. Do not use on softball fields. Do not use on tennis courts. Do not use on soccer fields. Do not use on football fields. Do not use on baseball fields. Do not use on softball fields.

8802

ProTurf

monocot weed control

8802

*Scotts*

## ProTurf

# monocot weed control

Selectively controls the following monocots:

|               |                   |
|---------------|-------------------|
| burnyardgrass | goosegrass        |
| crabgrass     | (in tiller stage) |
| dallisgrass   | parrotgrass       |
| (see below)   | (see below)       |
| foxtail       | witchgrass        |

**Directions for use**

Only weeds growing at the time of application will be affected. Weed seeds in the soil are not affected. Rain, mowing or heavy traffic within 24 hours reduces effectiveness by removing granules and may require a follow-up treatment. Apply to moist foliage at the normal rate any time susceptible weeds are in active growth. Localized infestations may be spot or patch treated. Two treatments 5 to 7 days apart are usually sufficient to achieve control. Repeat as needed. To quickly clear up crabgrass in the fall make two double rate applications. Dallisgrass and other paspalums are particularly difficult to control in California where especially favorable growing conditions exist.

**USE THESE SPREADER SETTINGS to ensure proper distribution**

25 . LBS TREATS 11,000 SQ FT ( . ACRE) AT NORMAL RATE  
 25 . LBS TREATS 5,500 SQ FT ( . ACRE) AT DOUBLE RATE

| SPREADER           | GROUND OR PTO SPEED | WIDTH OF COVERAGE | SPREADER RATE | SPREADER SETTING |
|--------------------|---------------------|-------------------|---------------|------------------|
| Scotts (Drop Type) | 3 mph               | overlap wheels    | 5             | 7                |
| Scotts 1000-B R    | 4.5 mph             | overlap wheels    | 5             | 7                |
| Gandy 8 R A Series | 4.5 mph             | overlap wheels    | 19            | 26               |
| 42 model           | 2.5 mph             | overlap wheels    | 34            | 44               |

**Precautions**

Do not use on lawns of St. Augustine, centipede, bahia or carpetgrass except as a non-selective spot treatment. Delay planting grass seed until after a heavy rain or watering and not sooner than 48 hours after last application. Delay treatment to new grass seedlings until after the new grass has been mowed 3 times. In hot weather, desirable turfgrasses and dichondra may brown temporarily, but will recover with watering.

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040  
 Oakland, Calif. 94604

8802

ACCEPTED  
 Feb. 26, 1978

537-85