


TREATS 5,000 SQ FT



weed control
with fertilizer

Lawn Pro

SCOTT'S BEST LAWN PRODUCTS

~~weed control~~



~~weed control~~

10-5-3

Fertilizes grass, controls dandelions
and other weeds (see list on back)

NET WEIGHT 14.7 LBS (5.0 KG)


For use on lawns, golf courses, and other areas where grass is grown. Apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall.



~~weed control~~

~~weed control~~


10-5-3



For use on lawns, golf courses, and other areas where grass is grown. Apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall.

For use on lawns, golf courses, and other areas where grass is grown. Apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall. For best results, apply to the lawn in the spring and fall.

Set control with fertilizer



30-5-3

Fertilizes grass, controls these rosette and vining weeds:

aster	evening	pepperweed
bitterress	primrose	pigweed
black medic	finch	plantain
blackhorn	johnsoga	prickly lettuce
butonweed	vine and ivy	puncturevine
chickweed	rice all	purslane
chicory	horset	ragweed
chickweed	knottweed	shepherds
clover	(seedling)	purse
(white)	lambquarters	smartweed
cutweed	lipid	stitchwort
curly dock	(matchweed)	thick leaved
daisy	monkswart	sandwort
floribane	morning glory	velvet
dandelion	mustard	whitegrass
English daisy	pennywort	yellow rocket

Use this spreader setting
to insure proper distribution

1 lb. treats 5,000 sq. ft.
Scotts Standard or Automatic Model... 5
(overlap wheels)

FOR USE ON TURF GRASS AREAS OTHER THAN PUTTING GREENS

Use when weeds are growing profusely. Growth should not
be so tall as to touch bottom of spreader. Apply to moist
forage. Repeat in one month if needed.

Rain, sprinkling, mowing or heavy traffic within 24 hours
after application knocks granules off weeds, reduces ef-
fectiveness.

Re-treating may be necessary, but wait at least 3 weeks to
observe full results of first application.

PRECAUTIONS

Avoid application on garden plants, flowers and shrubs.

On new plantings delay application until after the fourth
mowing.

Avoid application when daytime temperatures are consist-
ently below 60° F.

Delay sowing grass seed until after a heavy rain or water-
ing and not later than 2 months after application.

Not for use on creeping bentgrass, St. Augustine carpet-
grass, bermudagrass and other ground covers.

O. M. Scott & Sons

Marysville, Ohio 43040

XXXXXX Cameraman's Note
Poor Copy