

# NEW HIGH DENSITY fertilizer *plus* DSB fungicide



**NEW**  
HIGH DENSITY

## 28-0-7

# fertilizer *plus* DSB fungicide

FOR USE ON TURFGRASSES ONLY

Prevents or arrests fungus activity causing dollar spot, brownpatch and fusarium patch (*Fusarium nivale*) diseases. Fertilizes grass.

### USE THESE SPREADER SETTINGS to ensure proper distribution

35 Lbs. (15.9 Kg) treats 22,000 sq ft (1/4 Acre/2045 sq m) at NORMAL RATE  
35 Lbs. (15.9 Kg) treats 11,000 sq ft (1/4 Acre/1022 sq m) at DOUBLE RATE

| SPREADER                 | GROUND OR<br>PTO SPEED | WIDTH OF<br>COVERAGE | SPREADER SETTING<br>NORMAL<br>RATE | DOUBLE<br>RATE |
|--------------------------|------------------------|----------------------|------------------------------------|----------------|
| Scotts (Drop Type)       | 3 mph                  | overlap wheels       | 3 1/2                              | 4 3/4          |
| Scotts Rotary            | 3 mph                  | 10 feet              | D                                  | G              |
| Lely-WTR &<br>WFR Models | 4.5 mph                | 24 feet              | 2 1/2 III                          | 3 1/4 III      |
| Lely-HR<br>(PTO Model)   | 450 rpm<br>@ 4.5 mph   | 24 feet              | 2 1/2 III                          | 3 1/4 III      |
| Gandy 8 ft A Series      | 4.5 mph                | overlap wheels       | 10                                 | 14             |
| 42" model                | 2.5 mph                | overlap wheels       | 24                                 | 28             |

### FOR USE ON BENTGRASS PUTTING GREENS AND OTHER TURFGRASS AREAS

Apply to either moist or dry foliage.

**DOLLARSPOT** (*Sclerotinia homoeocarpa*) and **BROWN-PATCH** (*Rhizoctonia solani*)—to prevent fungus activity treat in late spring at **NORMAL RATE**, repeat at 2 or 3 week intervals.

During prolonged periods of high temperature and humidity, the above intervals may have to be shortened for brownpatch prevention.

To control unexpected attacks of either fungus, treat immediately when damage is first noticed. Use **DOUBLE RATE** for dollarspot and **NORMAL RATE** for brownpatch. Follow with preventative program outlined above.

\*If treatment is not effective following use of this product as recommended, a tolerant strain of fungi may be present. Contact your ProTurf Tech Rep. consideration should be

Prevents or arrests fungus activity causing dollarspot, brownpatch and fusarium patch (*Fusarium nivale*) diseases. Fertilizes grass.

# USE THESE SPREADER SETTINGS to ensure proper distribution

35 Lbs. (15.9 Kg) treats 22,000 sq ft (1/4 Acre/2048 sq m) at NORMAL RATE  
35 Lbs. (15.9 Kg) treats 11,000 sq ft (1/4 Acre/1022 sq m) at DOUBLE RATE

| SPREADER                         | GROUND OR<br>PTO SPEED | WIDTH OF<br>COVERAGE             | SPREADER SETTING<br>NORMAL<br>RATE | SPREADER SETTING<br>DOUBLE<br>RATE |
|----------------------------------|------------------------|----------------------------------|------------------------------------|------------------------------------|
| Scotts (Drop Type)               | 3 mph                  | overlap wheels                   | 3 1/2                              | 4 3/4                              |
| Scotts Rotary                    | 3 mph                  | 10 feet                          | D                                  | G                                  |
| Lely-WTR &<br>WFR Models         | 4.5 mph                | 24 feet                          | 2 1/2 III                          | 3 1/4 III                          |
| Lely-HR<br>(PTO Model)           | 450 rpm<br>@ 4.5 mph   | 24 feet                          | 2 1/2 III                          | 3 1/4 III                          |
| Gandy 8 ft A Series<br>42" model | 4.5 mph<br>2.5 mph     | overlap wheels<br>overlap wheels | 10<br>24                           | 14<br>28                           |

## FOR USE ON BENTGRASS PUTTING GREENS AND OTHER TURFGRASS AREAS

Apply to either moist or dry foliage.

**DOLLARSPOT\*** (*Sclerotinia homoeocarpa*) and **BROWN-PATCH** (*Rhizoctonia solani*)—to prevent fungus activity treat in late spring at **NORMAL RATE**, repeat at 2 or 3 week intervals.

During prolonged periods of high temperature and humidity, the above intervals may have to be shortened for brownpatch prevention.

To control unexpected attacks of either fungus, treat immediately when damage is first noticed. Use **DOUBLE RATE** for dollarspot and **NORMAL RATE** for brownpatch. Follow with preventative program outlined above.


\*If treatment is not effective following use of this product as recommended, a tolerant strain of fungi may be present. Contact your ProTurf Tech Rep, consideration should be given to prompt use of other suitable ProTurf fungicides.

**FUSARIUM PATCH** (*Fusarium nivale*) — to control fungus activity, apply at **DOUBLE RATE** at first sign of disease. Repeat if needed.

ProTurf Division, OM Scott & Sons, Marysville, Ohio 43040

## STORE THIS PRODUCT IN A COOL PLACE

### HOW TO STORE OPENED BAGS



1. Roll down top to seal out moist air.



2. Tie as shown to maintain seal.

ACCEPTED WITH COMMENTS

NET WT 24 LBS

# NUTRITIONAL SPRAY T-4 FOR CITRUS

## DIRECTIONS FOR USE FOR NUTRITIONAL DEFICIENCIES

Swift Nutritional Spray T-4 contains basic copper sulfate, neutral zinc, and neutral manganese. Use at rate of 4.8 pounds per 100 gallons of water. When used at recommended rate the standard dosage of copper, zinc, and manganese will be supplied. For best results, apply as a dormant or post-bloom spray.

## FOR FUNGUS DISEASES OF CITRUS

Melanose and Scab—Use at rate of 4.8 pounds per 100 gallons of water, one to three weeks after petal fall. Where melanose has been severe in the past or during wet springs, a second application should be made 4 weeks later. On varieties severely affected by scale make one application prior to spring flush of growth and another one to three weeks after petal fall.

## CONTAINS

## COPPER—ZINC—MANGANESE.

### ACTIVE INGREDIENTS:

Copper (in basic copper sulfate)  
as metallic ..... 15.60%

INERT INGREDIENTS: ..... 84.40%

Total 100.00%

### SECONDARY PLANT FOODS:

Zinc as ZnO ..... 26.00%

Manganese as MnO ..... 20.20%

E.P.A. Reg. No. 557-1871

E.P.A. Est. No. 557-FL-1

**CAUTION:** Keep Out Of Reach Of Children  
See Side Panel For Additional Cautions

## MIXING

Add to spray tank while running. It is not necessary for neutralization.

Harmful if swallowed. Wash with soap and water after use and foodstuffs.

ANTIDOTE: Call physician in case of poisoning. Treat with whites of egg.

This product is toxic to fish or ponds. Do not contaminate water, or disposal of waste specified on this label.

Do not reuse empty container. Do not waste or burning. Store in original container.

Swift Agricultural Chemicals Co.

NET WT 24 LBS

# NUTRITIONAL SPRAY T-4 FOR CITRUS

## CONTAINS COPPER – ZINC – MANGANESE.

### ACTIVE INGREDIENTS:

Copper (in basic copper sulfate)  
as metallic ..... 15.60%

INERT INGREDIENTS: ..... 84.40%  
Total 100.00%

### SECONDARY PLANT FOODS:

Zinc as ZnO ..... 26.00%  
Manganese as MnO ..... 20.20%

E.P.A. Reg. No. 557-1871

E.P.A. Est. No. 557-FL-1

**CAUTION:** Keep Out Of Reach Of Children  
See Side Panel For Additional Cautions

### MIXING INSTRUCTIONS

Add to spray tank when  $\frac{2}{3}$  full of water with agitator running. It is not necessary to add lime to this mixture for neutralization.

### CAUTION

Harmful if swallowed. Wash hands, arms, and face with soap and water after using. Avoid contamination of feed and foodstuffs.

ANTIDOTE: Call physician immediately. Give large quantities of whites of egg or milk.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burning with waste or burning. Stay away from smoke or fumes.

Manufactured By  
Swift Agricultural Chemicals Corporation  
(An Estech Company)  
Chicago, Illinois 60604