NEW CONCENTRATED FORMULA

Kwit chinchbug control

Treats 2,500 sq ft / $6.25

6433

Scotts No-Quibble Guarantee

If for any reason you are not satisfied with results after applying this product with a Scotts Spreader, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

NEW CONCENTRATED FORMULA

Scotts

Kwit®

CHINCHBUG CONTROL

controls chinchbugs

sod webworms (lawn moth larvae), cutworms, earwigs

A PROBLEM-SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE

NEW CONCENTRATED FORMULA

Kwit chinchbug control

NEW CONCENTRATED FORMULA

Kwit chinchbug control

NEW CONCENTRATED FORMULA

NEW CONCENTRATED FORMULA

6433

NET WEIGHT 5½ LBS.

6433

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. May be absorbed through the skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after using. Avoid contamination of food and foodstuffs. As directed, water lawn after application and allow to dry before permitting children or pets on treated areas. Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes

Apply this product only as specified on this label. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Active ingredient: Chlorpyrifos [0, 0-diethyl 0-(3,5,6 trichloro-2-pyridyl) phosphorothioate]	1.35%
Inert ingredients	98.65%
Total	100.00%

US Patent No 3,083,089 EPA Reg No 538-69 Product of USA

O M Scott & Sons Marysville, Ohio 43040

Treats 2,500 sq ft / NET WEIGHT 5½ LBS / $6.25

NEW CONCENTRATED FORMULA

Scotts

Kwit®

CHINCHBUG CONTROL

controls chinchbugs

sod webworms (lawn moth larvae), cutworms, earwigs

A PROBLEM-SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE

Directions for use

Apply over entire lawn area. Overlap spreader wheel tracks so no strips are missed. Water after application. Do not mow for 24 hours.

Before permitting children or pets on treated area, let lawn foliage dry off after watering in.

To control chinchbug—use anytime activity or damage is noticed (yellowing, then browning of grass often developing in spots then spreading through entire lawn.) Water thoroughly after application to wash particles into zone of insect activity.

To prevent chinchbug damage start applications as soon as daily temperatures reach 80 or above and repeat at two-month intervals.

To control sod webworms and cutworms—apply whenever their activity is observed or suspected. Sprinkle lightly after application to remove particles from leaves.

Use a Scotts Spreader (Standard or Automatic) at one of these settings to ensure proper distribution

NORMAL RATE—for chinchbug control (treats 2,500 sq ft) . . 7

LIGHT RATE—for other insects (treats 5,000 sq ft) 5½

♦ ♦ ♦

If you would like a free two-year subscription to our seasonal bulletin LAWN CARE, please write SCOTTS, Dept. 60, Marysville, Ohio 43040.

NEW CONCENTRATED FORMULA

Kwit chinchbug control

6433

538-69

7/27/197

2

ACCEPTED

JUL 12 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 538-69 SUBJECT TO ATTACHED COMMENTS.