

FRONT:

1986 PACKAGE COPY FOR WESTERN  
LAWN INSECT CONTROL XXXXX

(Scotts) Western Lawn  
Insect Control

(NOTE: Above description will be inverted.)

Treats 5,000 sq ft (465 m<sup>2</sup>)

SCOTTS POLICY

No-Quibble

Guarantee

SEE BACK FOR DETAILS

(Scotts(R))

Western Lawn  
Insect Control

JAN 9 1937

Controls chinch bugs, sod webworms (lawn moth larvae), cutworms and other insects listed below.

ants  
armyworms  
billbug adults  
brown dog ticks  
crickets  
earwigs

European crane fly  
larvae (leatherjacket)  
flea beetles  
grasshoppers  
vegetable weevils  
(in dichondra)

This product is not recommended for control of subsurface insects such as grubs or mole crickets.

For use on grass or dichondra lawns.

NET WEIGHT 7.5 LBS (3.40 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

LEFT COLUMN

Hazards to Humans and Domestic Animals: CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. May be absorbed through the skin. Do not apply by hand. Wash thoroughly with soap and water after using. Wash contaminated clothing before reuse. Avoid contamination of feed or foodstuffs. As directed, water after application and allow to dry before permitting children or pets on treated areas. Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is antidotal.

FRONT CONTINUED:

Statement of Practical Treatment:

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

Environmental Hazards: This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

RIGHT COLUMN

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENT:

Chlorpyrifos [O, O-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] . . . . .	1.35%
INERT INGREDIENTS: . . . . .	98.65%
Total	100.00%

EPA Reg No 538-69

EPA Est No 538-OH-1

Product of USA

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) XXXX, The O M Scott & Sons Company. All rights reserved.

RIGHT GUSSET

(Scotts(R)) Western Lawn Insect Control

XXXXX

SCOTTS POLICY  
No-Quibble  
Guarantee  
SEE BACK FOR DETAILS

LEFT GUSSET

XXXXX

(Scotts(R)) Western Lawn Insect Control

SCOTTS POLICY  
No-Quibble  
Guarantee  
SEE BACK FOR DETAILS

BACK:

EASY  
OPEN

PULL TAB  
and remove for  
pour spout

(Scotts(R))

Treats 5,000 sq ft (465 m<sup>2</sup>)

Western Lawn  
Insect Control

WLIC XXXXX  
Proof of  
Purchase  
Scotts

controls these insects:

sod webworms      chinch bugs      cutworms  
(lawn moth larvae)

ants, armyworms, billbug adults, brown dog ticks, crickets,  
earwigs, flea beetles, grasshoppers, European crane fly  
larvae (leather-  
jacket) vegetable weevils (in dichondra).

for use on grass or dichondra lawns

EASY TO APPLY

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- Apply when insect activity or damage (chewed or brown grass blades) is first observed in the lawn.
- Apply when lawn is dry.
- Water lawn after application to wash granules from the lawn foliage. If treating for chinch bugs, billbug adults or crane fly larvae, water thoroughly (1/4 inch) so active ingredient will be carried into the root zone. Don't allow children or pets on the lawn until it has dried.
- Treat the entire lawn to prevent further insect damage.
- You should not use any other lawn control product for at least one week after application.
- A second application may be needed in 6 weeks if insect activity continues.
- Do not use on fruit trees or vegetable plants.

BACK (continued):

Application Rate	Scotts Drop Spreaders	Scotts EasyGreen™ Rotary
For sod webworms, armyworms flea beetles, earwigs, vegetable weevils and cutworms (treats 5,000 sq ft)	X	X
For chinch bugs (also ants, billbug adults, European crane fly larvae, ticks, crickets and grasshoppers) (treats 3,500 sq ft)	X	X

CHINCH BUGS -- Start treatment when daily high temperatures are 80° or above. A second application may be needed in 6 weeks if insect activity continues.

BILLBUG ADULTS -- Apply in the spring (normally April-May) when adults emerge from over-wintering and before egg laying begins.

EUROPEAN CRANE FLY LARVAE (leatherjackets) -- Apply when activity is first noticed in the spring, usually mid-March to mid-April.

GRUBS and MOLE CRICKETS -- This product is not recommended for control of these or other subsurface insects.

TIPS ON SPREADING WITH SCOTTS SPREADERS

Scotts Drop:

- First apply 2 header strips
- Operate spreader the longest way of the lawn
- Overlap wheels to insure complete insect control
- Shut off spreader when you have reached the header trips or anytime you stop
- Mow lawn first if grass is tall enough to touch the bottom of spreader

EasyGreen™ Rotary:

- Slide shut-off lever to the right. Application automatically starts and stops when you do
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs

(ILLUS.)

(ILLUS.)

BACK (continued):

YOUR SATISFACTION GUARANTEED

XXXXX

Scotts Policy  
No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTTS...*the lawn people*  
MARYSVILLE, OHIO 43041

0 (UPC)

32247 XXXXX

(NOTE: The description below will be inverted.)

XXXXX

(Scotts) Western Lawn  
Insect Control