

Scotts
Shrub & Tree
WEED PREVENTER
plus FERTILIZER...



Shrub & Tree

WEED PREVENTER

plus FERTILIZER 20-4-8

- ✓ prevents sprouting of 26 types of weeds and grasses - see list on other side.
- ✓ prolonged feeding action - for deciduous and evergreen shrubs, trees and ground covers.



Net Weight 20 lbs

CAUTION - KEEP OUT OF REACH OF CHILDREN
 Hazardous if swallowed or inhaled. Avoid contact with skin, eyes, nose, or mouth.
 Directions: For use on shrubs, trees, and ground covers. Apply to soil around plants.
 Net Weight: 20 lbs. Scott's Shrub & Tree Weed Preventer, Fertilizer, and Fungicide.
 Active ingredient: Trifluralin (100% active ingredient).
 EPA Reg. No. 135-100001-1.
 For more information, contact Scotts Company, P.O. Box 130, Kenilworth, N.J. 07033.
 ©1985 Scotts Company.

5033

Scotts
Shrub & Tree
WEED PREVENTER
plus FERTILIZER...

5033

Scotts
Shrub & Tree
WEED PREVENTER
plus FERTILIZER...

NET WEIGHT 20 LBS / \$6.95

Scotts
Shrub & Tree
WEED PREVENTER
plus FERTILIZER...

5033

NET WEIGHT 20 LBS / \$6.95

Scotts
Shrub & Tree
WEED PREVENTER
plus FERTILIZER...

Broadleaf weeds prevented	Annual grasses prevented
carpetweed	banysardens
chickweed	brachiaria
Florida purslane (puslet)	bromegrass
goosefoot	cheat
knottweed	crabgrass
koehia (Mexican fireweed)	foxtail
lambquarter	gossypias
pigweeds	Johnsongrass (from seed)
purslane	Poa annua
tinging-nettle	sandbar
	stinkgrass
	panicum
	wild oats
	bahia
	crabgrass



Directions for use

In Zone A apply in the spring and again 3 to 6 months later. In Zone B add a third treatment in mid fall for best ground benefits.

Apply to soil around plants. Do not apply to lawn areas, or vegetables, or potted plants.



Drugstore diameter	Area Sq. Ft.	Amount Required
12"	1.1	1/2 cup
18"	2.2	1 cup
24"	3.3	1 1/2 cups
30"	4.4	2 cups
36"	5.5	2 1/2 cups
42"	6.6	3 cups
48"	7.7	3 1/2 cups
54"	8.8	4 cups
60"	9.9	4 1/2 cups

Maximum application rate: 4 lbs. per 1,000 sq. ft. of soil.
 Do not apply to lawn areas, or vegetables, or potted plants.
 Do not use when planting flowers from seed.

Scotts
No-Quibble
Guarantee

5033

5033

Net Weight 20 lbs

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. In cases of contact, flush with water.

Direct contamination of any body of water with the granular material may kill fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes. Do not reuse container. Destroy when empty. Do not contaminate feed or foodstuffs.

Net Weight 20 lbs Shrub & Tree Weed Preventer plus Fertilizer 20-4-8 guaranteed analysis*

Active ingredient: Trifluralin (*a, a, a*-trifluoro-2,6-dinitro-N,N-dipropyl-*p*-toluidine) 0.68%

Inert ingredients: Relative to herbicidal activity. 99.32%

Total 100.00%

Total nitrogen 20.00%

1.0% ammoniacal nitrogen from ammoniated phosphate

12.3% water soluble nitrogen from urea and methylene ureas

6.7% water insoluble nitrogen from methylene ureas

Available phosphoric acid (P₂O₅) from ammoniated phosphate 4.00%

Soluble potash (K₂O) from muriate of potash 8.00%

Chlorine 6.25%

Potential acidity equivalent to 500 lbs calcium carbonate per ton

*In Michigan reads "minimum guaranteed analysis"

US Pat No 3,076,700 EPA Reg No 538-83-AA Product of USA

O M Scott & Sons Marysville, Ohio 43040

5033

5033



O & TEE

**WEED PREVENTER
plus FERTILIZER**
20-4-8

UNDER THE FERTILIZER...
MUNICIPAL...
TOWN...
REGISTERED

bigweeds
purslane
tinging nettle

Johnsongrass
Poa annua
sandbur



Directions for use

In Zone A apply in the spring and again 7 to 8 months later. In Zone B add a third treatment in mid-fall for year-round benefits.

Use on bare soil only. Do not use for trees or shrubs growing in the lawn.



Cultivate soil first to loosen it, and to eliminate existing weeds. Then make application by scattering material evenly at the rate of 4-1/2 cups (equivalent to a one quart container, or a one-pound coffee can) per 10' x 10' area.



If applying to individual trees, shrubs or plants, scatter the material evenly beneath the spread of the branches (the dripzone) as follows:

<u>Dripzone diameter</u>	<u>Area (Sq ft)</u>	<u>Amount required</u>
1 ft	3-4	1-2 tablespoon
2 ft	3	2 tablespoon
3 ft	7	4-1/2 tablespoon
4 ft	13	1-2 cup
5 ft	20	1 cup
10 ft	78	3-3/4 cups

Rake material lightly into the top inch of soil. Then water the area evenly, and sufficiently to wet the top inch of soil, and to wash material off the leaves of plants.

May be applied to large areas of bare soil with a Scotts Spreader (Standard or Automatic model) using setting #8-1-2. Do not apply with a rotary type spreader.

When transplanting annual and perennial flowers, wait at least one week after moving plant before using this product.

Decorative mulches can be used after application but the soil should not be disturbed.

**Do not apply to bulb plants, or to ajuga, dichondra,
pachysandra, or myrtle, or to lawn areas, or vegetables, or potted plants.
Do not use when planting flowers from seed.**



5033

Scotts

If for any reason you are not satisfied with results after using this prod-

Scotts

**WEED PREVENTER
plus FERTILIZER**
20-4-8



Directions for use

In Zone A apply in the spring, and again 2 to 3 months later. In Zone B, add a third treatment in mid-fall for year-round benefits.

Use on bare soil only, do not use for trees or shrubs growing in the lawn.



Cultivate soil first to loosen it, and to eliminate existing weeds. Then make application by scattering material evenly at the rate of 4-1/2 cups (equivalent to a one quart container or a one-pound coffee can) per 10 x 10 area.



If applying to individual trees, shrubs or plants, scatter the material evenly beneath the spread of the branches (the dripzone) as follows:

Dripzone diameter	Area (Sq ft)	Amount required
1 ft	3.4	1/2 tablespoon
2 ft	3	2 tablespoon
3 ft	7	4-1/2 tablespoon
4 ft	13	1/2 cup
5 ft	20	1 cup
10 ft	78	3-3/4 cups

Rake material lightly into the top inch of soil. Then water the area evenly, and sufficiently to wet the top inch of soil, and to wash material off the leaves of plants.

May be applied to large areas of bare soil with a Scotts Spreader (Standard or Automatic model) using setting = 8-1/2. Do not apply with a rotary type spreader.

When transplanting, annual and perennial flowers, wait at least one week after moving plant before using this product.

Decorative mulches can be used after application but the soil should not be disturbed.

Do not apply to bulb plants, or to ajuga, dichondra, pachysandra, or myrtle, or to lawn areas, or vegetables, or potted plants. Do not use when planting flowers from seed.

5033

**Scotts
No-Quibble
Guarantee**

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTT'S the lawn people
Morton, IL 61704
1-800-333-3333