

PLEASE NOTE

This image contains more than one label approved for this product on this date.

Best Available Copy

4214

Treats 5,000 sq ft

Scott's

Lawn Insect Control plus Fertilizer

Treats 5,000 sq ft

Scott's

Lawn Insect Control plus Fertilizer

✓controls insects ✓feeds the lawn

Controls white grubs, Japanese beetle, European chafer, Southern chafer, *Phyllophaga* (May or June beetle).

Also controls the following insects on grass or herbaceous lawns:
army worms, chinchbugs, cutworms, leafhoppers, Bermuda mites, clover mites, flies, male crickets, billbugs, crickets, fire beetles (in Florida), chiggers, curculionids (in Dickcissel), ticks, and webworms (also moth larvae) vegetable weevils

A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE

How To Use

- DO apply when insect activity or damage (chewed or browned leaves) is first observed in the lawn. For grubs, the most likely period is from late July through early October; the most likely period is Spring. For mole crickets, the best time to apply is in May or June. DO apply when lawn is dry. DO water lawn after application to wash granules from the lawn foliage. If treating for grubs, water thoroughly so that active ingredients will be carried into the root zone. DO make a repeat application of Scott's Lawn Insect Control with or without fertilizer if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 8 weeks.)
- DO apply at Scott's Spreader (Standard or Automatic Model) setting 5 (Treats 5,000 sq ft)
- DO NOT treat just the browned areas of lawn. To prevent further damage, treat entire lawn.
- DO NOT use any other lawn control product for at least one week after application.
- DO NOT allow traffic on lawn after application until granules are worked off lawn foliage and lawn has dried.
- DO NOT use on fruit or vegetable plantings.
- DO NOT apply with a rotary-type spreader.

Free bulletin-service

Form with fields for name, address, and phone number for free bulletin service.

Scott's No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send an unopened jar back and we will issue you a refund. Check promptly. SCOTT'S, The Lawn People, MARYSVILLE, OHIO 43040

A SCOTT'S LAWN PRO. PRODUCT. Includes 1/2 lb. of available lawn fertilizer.

Scott's

Lawn Insect Control plus Fertilizer

- ✓controls white grubs, sod webworms, chinchbugs- and other lawn insects listed on back panel.
- ✓gives your lawn a full feeding of Scott's long lasting fertilizer

A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE

Lawn Insect Control plus Fertilizer

Scott's

NET WEIGHT 15 1/2 LBS (treats 5,000 sq ft)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. May be absorbed through skin. As directed water lawn after application and allow to dry before permitting children and pets on treated area.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

Active ingredient: 0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 3.60% 96.40%
Inert ingredients: 96.40%
EPA Reg. No. 538-92 EPA Est. 574-OH-1 Product of U.S.A. Total 100.00%
Net Weight 15 1/2 lbs. Lawn Insect Control Plus Fertilizer 28-6-4 guaranteed analysis
Total nitrogen (N) 28%
1.3% ammoniacal nitrogen from ammoniated phosphate
17.3% water soluble nitrogen from urea and methylene ureas
9.4% water insoluble nitrogen from methylene ureas
Available phosphoric acid (P2O5) from ammoniated phosphate 6%
Soluble potash (K2O) from potassium sulfate 4%
Potential acidity equivalent to 1200 lbs calcium carbonate per ton.
U.S. Pat. Nos. 3,231,363; 3,705,794 Other Pat. Pend. O.M. Scott & Sons, Marysville, Ohio 43040

4214

ACCEPTED
MAY 18 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 538-92
EPA Reg. No. 538-92

4214

Scott's

Lawn Insect Control plus Fertilizer

538-92

5-18-76

1041

NEXT

LABEL

PLUS FERTILIZER

✓controls insects ✓feeds the lawn

Controls white grubs; Japanese beetle, European chafer, Southern chafer, *Phyllophaga* (May or June beetle).

Also controls the following insects in grass or dichondra lawns:

armyworms	chinchbugs	earwigs	leafhoppers
Bermuda mites	clover mites	fleas	mole crickets
billbugs	crickets	flea beetles	(in Florida)
chiggers	cutworms	(in Dichondra)	ticks
sod webworms (lawn moth larvae)	vegetable weevils		

A PROBLEM-SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE

lients will be carried into the root-zone).

DO make a repeat application of Scotts Lawn Insect Control (with or without fertilizer) if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 6 weeks.)

DO apply at Scotts Spreader (Standard or Automatic Models) setting

5

DON'T treat just the browned areas of lawn. To prevent further damage, treat entire lawn.

DON'T use any other lawn control product for at least one week after application.

DON'T allow traffic on lawn after application until granules are watered off lawn foliage and lawn has dried.

DON'T use on fruit or vegetable plantings.

DON'T apply with a rotary-type spreader.

NET WEIGHT 31 LBS (treats 10,000 sq ft)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. May be absorbed through skin. As directed water lawn after application and allow to dry before permitting children and pets on treated area.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

Active ingredient: 0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 3.60%
Inert ingredients: 96.40%

EPA Reg. No. 538-92	EPA Est 538-OH-1	Product of U S A	Total 100.00%
Net Weight 31 lbs	Lawn Insect Control Plus Fertilizer	28-6-4	guaranteed analysis
Total nitrogen (N)			28%

1.3% ammoniacal nitrogen from ammoniated phosphate	
17.3% water soluble nitrogen from urea and methylene ureas	
9.4% water insoluble nitrogen from methylene ureas	
Available phosphoric acid (P ₂ O ₅) from ammoniated phosphate	6%
Soluble potash (K ₂ O) from potassium sulfate	4%

Potential acidity equivalent to 1200 lbs calcium carbonate per ton.
US Pat Nos 3,231,363; 3,705,794 Other Pat Pend O M Scott & Sons, Marysville, Ohio 43040

86A4

