



Lawn Insect Control plus Fertilizer

✓controls insects ✓feeds the lawn

Controls white grubs; Japanese beetle, European chafer, Southern chafer, *Phyllophaga* (May or June beetle).

Also controls the following insects in grass or dichondra lawns:

armyworms	chinchbugs	earwigs	leafhoppers
Bermuda mites	clover mites	fleas	mole crickets
billbugs	crickets	flea beetles	(in Florida)
chiggers	cutworms	(in Dichondra)	ticks
soil webworms (lawn moth larvae)		vegetable weevils	

A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE.

How To Use:

- DO apply when insect activity or damage (weeched or browned leaves) is first observed in the lawn.
For grubs, the most likely period is from late July through early October, the next most likely period is Spring.
For mole crickets, the best time to apply is in May or June.
- DO apply when lawn is dry.
- DO water lawn after application to wash granules from the lawn foliage. If treating for grubs, water thoroughly (so that active ingredients will be carried into the root zone).
- DO make a repeat application of Scott's Lawn Insect Control (with or without fertilizer) if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 6 weeks.)
- DO apply at Scott's Spreader (Standard or Automatic Models) setting **5 1/2**.
(Treats 10,000 sq ft)
In Colorado use setting... **.6**
(Treats 8,000 sq ft)
- DO apply at Scott's Rotary-Type Spreader Setting... **F**
(Treats 10,000 sq ft)
- DON'T treat just the browned areas of lawn. To prevent further damage, treat entire lawn.
- DON'T use any other lawn control product for at least one week after application.
- DON'T allow traffic on lawn after application until granules are watered off lawn foliage and lawn has dried.
- DON'T use on fruit or vegetable plantings.

NET WEIGHT 31 LBS (treats 10,000 sq ft)
(in Colorado treats 8,000 sq ft)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. May be absorbed through skin. As directed water lawn after application and allow to dry before permitting children and pets on treated area.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

Active ingredient: 0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 3.60%
Inert ingredients: 96.40%
EPA Reg. No. 538-92 EPA Est 538-OH-1 Product of U S A Total 100.00%
Net Weight 31 lbs Lawn Insect Control Plus Fertilizer 28-6-4 guaranteed analysis
Total nitrogen (N) 28%
1.3% ammoniacal nitrogen from ammoniated phosphate
17.3% water soluble nitrogen from urea and methylene ureas
9.4% water insoluble nitrogen from methylene ureas
Available phosphoric acid (P₂O₅) from ammoniated phosphate 6%
Soluble potash (K₂O) from potassium sulfate 4%
Potential acidity equivalent to 1200 lbs calcium carbonate per ton.
US Pat Nos 3,231,363; 3,705,794 Other Pat Pend O M Scott & Sons, Marysville, Ohio 43040

86A4

ACCEL TO
with 0-0-0-0-0-0
In 1-1-1-1-1-1

BEST DOCUMENT AVAILABLE

MADE IN U.S.A.
This product is registered under EPA Reg. No. 538-92

301/94672
17/5

Dr. Michael Kelty
O. M. Scott & Sons Company
Marysville, OH 43041

Dear Dr. Kelty:

Slaw

Subject: Amendment - Add Rotary Spreader Directions
Lawn Insect Control Plus Fertilizer
EPA Registration No. 538-92
Your Application Dated November 5, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Your label should be updated according to our January 24, 1983 letter.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

PM:LARocca:DCR-28758:WANG-0010C:hk:Raven:479-2013:4/27/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							



72% CHLORDANE

EMULSIFIABLE CONCENTRATE

For Professional Control of Termites

Only For Use and Storage by Commercial Pest Control Operators

ACTIVE INGREDIENTS:

Technical Clordane* 72%
Kerosene 20%

INERT INGREDIENTS: 8%
TOTAL: 100%

*Equivalent to 43.2% Octachloro-4,7-Methanotetrahydroindane and 28.8% related compounds.

ACCEPTED
APPLICANT'S
LABELING

MAY 5 1983

KEEP OUT OF THE REACH OF CHILDREN

WARNING

728-47

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center immediately.
Drink one or two glasses of water and induce vomiting by touching the
back of throat with finger. Repeat until vomit fluid is clear. Do not
induce vomiting or give anything by mouth to an unconscious person.
IF INHALED: Remove victim to fresh air. Apply artificial respiration
if indicated.
IF ON SKIN: Remove contaminated clothing and wash affected areas with
soap and water.
IF IN EYES: Flush eyes with plenty of water. Get medical attention
immediately.
See Side Panel for Additional Precautionary Statements.

EPA Reg. #728-47
EPA Est. #728-AL-1

NET CONTENTS: 1 gallon

Manufactured by: Southland Pearson & Company
P.O. Box 7344
Mobile, AL 36607