

Treats 5,000 sq ft (465 m²)



Lawn Insect Control plus Fertilizer



39032

ACCEPTED

JAN 16 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 538-92

Scott's Lawn Insect Control plus Fertilizer



Lawn Insect Control plus Fertilizer

Lawn Insect Control plus Fertilizer Scott's

For use on Lawn and Turfgrasses only

Controls white grubs, sod webworms, chinch bugs[™] and other lawn insects listed on back panel.

Gives your lawn a full feeding of Scott's long lasting fertilizer.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

A SCOTT'S **Lawn Pro** PRODUCT

Available only at authorized Lawn Pro stores

39032

NET WEIGHT 15½ LBS (7.03 kg)

28-6-4

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals: CAUTION. Harmful if swallowed. May be absorbed through the skin. Do not get on skin or clothing. Avoid inhalation. If on skin, remove contaminated clothing and wash affected area with soap and water. If on clothing, flush with plenty of water. Call a physician if irritation persists. Wash thoroughly after handling and before eating or smoking. Do not consume, use, feed or live with. Wash treated area after application and allow to dry before reentering use.
Signs and Symptoms: Dizziness in a cholinesterase inhibitor. Atropine is antidotal if symptoms of cholinesterase inhibition are present.

Environmental Hazards: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of a waste.
STORAGE: Store in a cool, dry place. Re-seal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: See separate container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse bag. Discard in trash.

ACTIVE INGREDIENT

Diazinon [10-Dithiopyr (2-ethyl-6-methylpyridin-3-ylmethyl) phosphorothioate] 1.00%
INERT INGREDIENTS: 99.00%
Total 100.00%

EPA Reg. No. 538-92 EPA Est. 530-011
Product of USA
Lawn Insect Control plus Fertilizer 28-6-4 643 0121
Controlled Analysis
Total nitrogen (N) 2.0%
1.2% ammoniacal nitrogen
1.2% urea, methylene urea, nitrogen
0.8% water insoluble nitrogen
Available phosphorus and (P₂O₅) 6.0%
Soluble phosphorus (P₂O₅) 4.0%
Derived from monoammonium phosphate, urea, methylene urea and potassium sulfate
US Pat. No. 3,303,794

(Other Pat. Pend.)

The D M Scott & Son Company
Marysville, Ohio 43041

39032

BEST AVAILABLE COPY

Easy
Open



Treats 5,000 sq ft (465 m²)



Scotts Lawn Insect Control plus Fertilizer

Controls insects and feeds the lawn.

Controls white grubs such as Japanese beetle, European chafer, Masked chafer, *Phyllophaga* (May or June Beetle.)

Also controls the following insects in grass or dichondra lawns:

ants	chinch bugs	European crane fly larvae (leatherjackets)	leafhoppers	ticks
armyworms	clover root crickets	fleas	mole crickets and webworms (lawn moth larvae)	vegetable weevil (in dichondra)
Bermuda nibbler	cutworms	fire beetles (in dichondra)		
chiggers	onions			

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO apply when insect activity or damage (chewed or browned leaves) is first observed in the lawn.

NOTE: For Grub Control

Northern States—The most likely period for grubs is late July through September but in some cases grubs are active in the spring. An application for spring grub activity requires a repeat summer-early fall application for effective control.

Southern States—Apply during July-August. A second application may be needed 4-6 weeks later. Applications made later in the year, when grubs are more mature, are less effective.

For chinch bugs start treatments when daily high temperatures are 80° or above and repeat at 6 week intervals.

For mole crickets, the best time to apply is in May to July.

DO apply when lawn is dry. For mole crickets and white grubs, water thoroughly, let lawn dry and then apply product.

DO water lawn after application to wash granules from the lawn foliage. If treating for grubs or mole crickets, water thoroughly (so that active ingredients will be carried into the rootzone).

DO make a repeat application of Scotts Lawn Insect Control (with or without fertilizer) if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 6 weeks.)

Application Rate	Scotts Drop Spreader	Scotts EasyGreen [™] Rotary
(Treats 5,000 sq ft)	6	26

DON'T treat just the browned areas of lawn. To prevent further damage, treat entire lawn.

DON'T use any other lawn control product for at least one week after application.

DON'T allow traffic on lawn after application until granules are watered off lawn foliage and lawn has dried.

DON'T use on fruit or vegetable plantings.

A SCOTT'S **Lawn Pro** PRODUCT
Available only at authorized Lawn Pro stores

Scotts Policy No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, we are sure to get your money back. Simply send us proof of purchase and we will send you a refund check promptly.

SCOTT'S ... for the people • Marysville, Ohio 43041

BEST AVAILABLE COPY



Scotts Lawn Insect Control plus Fertilizer



39032