



WESTERN Lawn Insect Control plus Fertilizer

✓ controls lawn moth larva (sod webworms),
flea beetles, cutworms

chinchbugs vegetable weevils crickets grasshoppers
ants brown dog ticks earwigs

✓ gives your lawn a feeding of Scotts long lasting fertilizer

For use on grass or dichondra lawns

A PROBLEM-SOLVING PRODUCT BY SCOTTS, THE LAWN PEOPLE

How To Use

DO use anytime activity or damage is noticed or suspected, to control any insect listed above.

DO apply at Scotts Spreader (Standard or Automatic Models) setting.... **5**
(Treats 15,000 sq ft)

In Colorado use setting.... **5½**
(Treats 12,900 sq ft)

When treating for flea beetles, sod webworms, vegetable weevils, earwigs, and cutworms

DO apply at Scotts Spreader (Standard or Automatic Model) setting. **7**
when treating for chinchbugs, ants, ticks, crickets and grasshoppers. At this setting, contents will treat 7,500 sq ft of lawn area.

DO overlapspreader wheel tracks so no strips are missed.

DO use as soon as daily temperatures reach 80° or above, to prevent chinchbug damage. Repeat treatment at two-month intervals.

DO water lawn thoroughly after application when treating for chinchbugs.

DO sprinkle lawn lightly after application when treating for all other controllable insects, to remove particles from grass blades.

DON'T spot treat damaged areas only—apply over entire lawn area to provide complete protection.

DON'T use when grass is wet and so tall it touches bottom of spreader (let lawn dry before making application).

DON'T mow for 24 hours after application.

DON'T allow children or pets on lawn area until grass has dried off (after watering in).

DON'T use any other control product (such as weed control) for one week after application.

ACCEPTED
JUN 7 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act
EPA Reg. No. 538-94

NET WEIGHT 43 LBS

30-5-3

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after using. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash contaminated clothing before re-use. Avoid contamination of food and foodstuffs. As directed, water lawn after application and allow to dry before permitting children or pets on treated areas. Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Active ingredient: Chlorpyrifos [0, 0-diethyl 0-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate] 0.55%

Inert ingredients: 99.45%

EPA Reg No 538-94-AA EPA Est. 538-OH-1 Product of USA Total 100.00%

Net Weight 43 lbs Western Lawn Insect Control plus Fertilizer 30-5-3 guaranteed analysis

Total nitrogen (N) 30%

1.2% ammoniacal nitrogen from ammoniated phosphate

18.8% water soluble nitrogen from urea and methylene ureas

10.0% water insoluble nitrogen from methylene ureas

Available phosphoric acid (P₂O₅) from ammoniated phosphate 5%

Soluble potash (K₂O) from potassium sulfate 3%

Potential acidity equivalent to 1200 lbs calcium carbonate per ton

US Process Pat Nos 3,231,363; 3,705,794 Other Pat Pend O M Scott & Sons, Marysville, Ohio 43040

86B3

Treats up to 15,000 sq ft




WESTERN Lawn Insect Control plus Fertilizer

- ✓ controls lawn moth larva (sod webworms),
flea beetles, cutworms
- chinchbugs vegetable weevils crickets grasshoppers
ants brown dog ticks earwigs
- ✓ gives your lawn a feeding of Scott's long lasting fertilizer

For use on grass or dichondra lawns
A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE

How To Use

- DO use anytime activity or damage is noticed or suspected, to control any insect listed above.
- DO apply at Scott's Spreader (Standard or Automatic Model) setting **5** when treating for flea beetles, sod webworms, vegetable weevils, earwigs and cutworms. At this setting, contents will treat 15,000 sq ft of lawn area.
- DO apply at Scott's Spreader (Standard or Automatic Model) setting **7** when treating for chinchbugs, ants, ticks, crickets and grasshoppers. At this setting, contents will treat 7,500 sq ft of lawn area.
- DO overlap spreader wheel tracks so no strips are missed.
- DO use as soon as daily temperatures reach 80° or above, to prevent chinchbug damage. Repeat treatment at two-month intervals.
- DO water lawn thoroughly after application when treating for chinchbugs.
- DO sprinkle lawn lightly after application when treating for all other controllable insects, to remove particles from grass blades.
- DON'T spot treat damaged areas only—apply over entire lawn area to provide complete protection.
- DON'T use when grass is wet and so tall it touches bottom of spreader (let lawn dry before making application).
- DON'T mow for 24 hours after application.
- DON'T allow children or pets on lawn area until grass has dried off (after watering in).
- DON'T use any other control product (such as weed control) for one week after application.



NET WEIGHT 43 LBS

30-5-3

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after using. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash contaminated clothing before re-use. Avoid contamination of food and foodstuffs. As directed, water lawn after application and allow to dry before permitting children or pets on treated areas. Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Active ingredient: Chlorpyrifos [0, 0-diethyl 0-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate] 0.55%
Inert ingredients: 99.45%

EPA Reg No 538-94-AA EPA Est. 538-OH-1 Product of USA Total 100.00%

Net Weight 43 lbs Western Lawn Insect Control plus Fertilizer 30-5-3 guaranteed analysis

Total nitrogen (N) 30%
1.2% ammoniacal nitrogen from ammoniated phosphate
18.8% water soluble nitrogen from urea and methylene ureas
10.0% water insoluble nitrogen from methylene ureas

Available phosphoric acid (P₂O₅) from ammoniated phosphate 5%
Soluble potash (K₂O) from potassium sulfate 3%

Potential acidity equivalent to 1200 lbs calcium carbonate per ton
US Process Pat Nos 3,231,363; 3,705,794 Other Pat Pend O M Scott & Sons, Marysville, Ohio 43040

86B3