

8797

8797

8797

NET WEIGHT 32 3/4 LBS (14.9 Kg)

ProTurf

NEW!
fungicide II

Scotts

ProTurf

NEW!
fungicide II

for prevention and control
of Pythium blight and gray snowmold

FOR PROFESSIONAL USE ONLY

NET WEIGHT 32 3/4 LBS (14.9 Kg)

8797

8797

ProTurf

NEW!
fungicide II

ACCEPTED

Feb. 25, 1974

538-103

8797



NEW!
fungicide II

prevents or arrests blighting activity of
gray snowmold and Pythium blight

USE THESE SPREADER SETTINGS
to insure proper distribution



FOR USE ON TURFGRASS AREAS

Whenever used on greens and fairways apply to moist foliage in the morning or evening. Sprinkle lightly just before traffic resumes to avoid pickup of material on shoes, golf balls or maintenance equipment.

Prevention of Pythium blight - apply when high temperatures and humidity favor development. Repeat at 5-7 day intervals as long as these weather conditions persist. Apply to moist foliage.

Control of Pythium blight - apply to moist foliage as soon as disease activity is observed. Repeat at 5-7 day intervals during high temperature and humidity conditions.

In case of severe disease attack make a double rate application to moist foliage and continue normal rate applications at 5-7 day intervals during high temperature and humidity conditions.

On new seedings apply at normal rate after sowing seed. At first signs of disease apply to moist foliage of newly germinated grass and repeat at 7-10 day intervals if evidence of disease persists.

When treating to control Pythium blight do not fertilize or water heavily until disease activity is arrested. Prolonged contact with foliage enhances effectiveness.

Gray snowmold - apply prior to first heavy snow early winter. Repeat in midwinter if turf is free of snow. A final treatment in late winter as the snow melts extends protection. Applications can be to dry or moist grass.

In areas where snow coverage is present the entire winter make a double rate application prior to the first snow and a normal rate application when snow melts in the spring.

ProTurf Division, O.M. Scott & Sons, Marysville, Ohio 43040

8797

NEW!

fungicide II

for prevention and control
of Pythium blight and gray snowmold

FOR PROFESSIONAL USE ONLY

NET WEIGHT 32 3/4 LBS (14.9 Kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN

May irritate eyes, nose, throat, and skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Do not reuse container, destroy when empty. Do not graze or feed clippings from treated areas to livestock.

Do not contaminate water by cleaning of equipment or disposal of water. Apply this product only as specified on this label.

Active ingredient: Chlorothal (1,4-dichloro-2,5-dimethoxybenzene) 6.25%
Inert ingredients 93.75%

EPA Reg. No. 538-103-AA
ProTurf Division, O.M. Scott & Sons

Total 100.00%
Product of USA
Mansfield, Ohio 43040

8797

fungicide II

ACCEPTED

Feb. 25, 1974

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTER-
ED UNDER NO. **538-103**

8797

Whenever use morning or evening avoid pickup equipment.

Prevention of humidity favors these weather

Control of Pyth activity is ob temperature and

In case of severe moist foliage intervals during

On new seed signs of disease and repeat at

When treating heavily until foliage enhanced

Gray snowmold repeat in midwinter as the to dry or moist

In areas where double rate application w

ProTurf Division

8797

fungicide II

NEW!
fungicide II

fungicide II

prevents or arrests damaging activity
of gray snowmold and Pythium blight

**USE THESE SPREADER SETTINGS
to ensure proper distribution**

32 1/2 LBS TREATS 11,000 SQ FT (1/3 ACRE) AT NORMAL RATE
32 1/2 LBS TREATS 5,500 SQ FT (1/6 ACRE) AT DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTING	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	6	8
Scotts Rotary	3 mph	8 feet	G	K
Lely-WTR & WFR Models	4.5 mph	16 feet	4 : III	6 III
Lely-HR (PTO Model)	450 rpm 4.5 mph	16 feet	4 : III	6 III
Gandy 8 ft A Series	4.5 mph	overlap wheels	18	24
42 Model	2.5 mph	overlap wheels	35	42

FOR USE ON TURFGRASS AREAS

Whenever used on greens and fairways apply to moist foliage in the morning or evening. Sprinkle lightly just before traffic resumes to avoid pickup of material on shoes, golf balls, or maintenance equipment.

Prevention of Pythium blight — apply when high temperatures and humidity favor development. Repeat at 5-7 day intervals as long as these weather conditions persist. Apply to moist foliage.

Control of Pythium blight — apply to moist foliage as soon as disease activity is observed; repeat at 5-7 day intervals during high temperature and humidity conditions.

In case of severe disease attack make a double rate application to moist foliage and continue normal rate applications at 5-7 day intervals during high temperature and humidity conditions.

On new seedings, apply at normal rate after sowing seed; at first signs of disease apply to moist foliage of newly germinated grass, and repeat at 7-10 day intervals if evidence of disease persists.

When treating to control Pythium blight do not fertilize or water heavily until disease activity is arrested; prolonged contact with foliage enhances effectiveness.

Gray snowmold — apply prior to first heavy snow (early winter). Repeat in midwinter if turf is free of snow. A final treatment in late winter as the snow melts, extends protection. Applications can be to dry or moist grass.

In areas where snow coverage is present the entire winter, make a double rate application prior to the first snow, and a normal rate application when snow melts in the spring.

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040

ACCEPTED

Feb. 25, 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR TOXIC SUBSTANCE CONTROL REGISTER-
ED UNDER NO. 538-103

8797