

# 13 control fertilizer

# 13 control fertilizer

USE ONLY  
CENTRAL TURFGRASS ONLY

H OF CHILDREN  
contact. In case of skin contact, wash  
h large quantities of water

pty container, destroy when empty  
pyny!) benzamide 32%  
99.68%  
Total 100.00%

Fertilizer 27-0-13 guaranteed analysis  
27%

0%  
0%  
18.0%  
9.0%  
0%  
0%  
13%  
40%  
0.12%

lfate  
onate per ton  
3-8623 Filed 1973 Product of USA  
Marysville, Ohio 43040  
Oakland, Calif. 94604

8623

ACCEPTED

April 11, 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 538-84

greens, apply at Normal Rate in the spring when weather favors bermudagrass regrowth.

For control of Kentucky bluegrass and bentgrass, in areas other than tees and greens, apply at Double Rate any time they are actively growing.

### USE THESE SPREADER SETTINGS to ensure proper distribution

37 1/2 LBS TREATS 11,000 SQ FT (1/4 ACRE) AT NORMAL RATE  
37 1/2 LBS TREATS 5,500 SQ FT (1/8 ACRE) AT DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTING NORMAL RATE	SPREADER SETTING DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	6	8
Scotts Rotary	3 mph	8 feet	H	K
Scotts 1000-8 ft	4.5 mph	overlap wheels	3'	6'
Lely-WTR & WFR Models	4.5 mph	18 feet	5 III	6' III
Lely-HR (PTO Model)	450 rpm @ 4.5 mph	18 feet	5 III	6' III
Gandy 8 ft A Series	4.5 mph	overlap wheels	15'	21
42' model	2.5 mph	overlap wheels	33	39

### Precautions

Do not use on dichondra, bluegrass, annual or perennial ryegrass, fescuegrasses, bentgrass or bahiagrass where these grasses are the desired turf.

Not for use on muck or peat soils.

Do not apply this product where winter overseeding with cool season turfgrasses such as bluegrasses, ryegrasses, bentgrasses or fescuegrasses is planned within 60 days.

Application on golf greens prior to a severe cold or freezing period can cause discoloration. Avoid winter applications where these conditions are likely to occur.

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040  
Oakland, Calif 94604

### PESTICIDE-FERTILIZER MIXTURE

CAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Avoid breathing dust, contact with skin or eyes. In case of skin contact, wash with soap and water. In case of eye contact, wash with large quantities of water. Avoid contamination of feed and food. Do not contaminate irrigation ditches or water used for irrigation purposes. Apply only as specified on this label. Do not reuse empty container, destroy when empty.

Contains 3.5 d-chloro N (1,1-dimethyl-2-propynyl) benzamide, 32% (6.4 lbs./ton), apply 37 1/2 lbs. per 11,000 sq ft at normal rate, 37 1/2 lbs. per 5,500 sq ft at double rate.


8623

3





**NEW**  
**FFII**  
Fertilizer and fungicide



853

# NEW 14-3-3 FFII

Fertilizer and fungicide for use on bluegrass, fescue, bluegrass-fescue mixtures, perennial ryegrass, fine fescue, bahia and centipedegrass.

Do not apply on bentgrasses, zoysia, or ricegrass.

Fertilizes grass. Prevents and controls stripe smut, leaf spot, rust, brownpatch, dollarspot, and snowmold. Increases and extends green color of grass.

### USE THESE SPREADER SETTINGS to provide proper distribution

35 LBS TREATS 11,000 SQ FT (1/4 ACRE) AT NORMAL RATE  
35 LBS TREATS 5,500 SQ FT (1/8 ACRE) AT DOUBLE RATE

SPREADER	SP. RATE	WHEEL OVERLAP	FEET	FEET
Scotts (Drop Type)	4 mph	overlap wheels	6	6
Scotts Rotary	3 mph	8 feet	5	5
Scotts 1000 8 ft	4.5 mph	overlap wheels	5	5
Lely WTR & WFR Models	4.5 mph	16 feet	5 IV	6 IV
Lely HR (PTO Model)	450 rpm = 4.5 mph	16 feet	5 IV	6 IV
Gandy 8 ft A Series 42 Model	4.5 mph	overlap wheels	19	26
	2.5 mph	overlap wheels	35	43

To prevent or control leafspot, dollarspot and snowmold. Apply at **NORMAL RATE**. Make applications in late fall and early spring to prevent leafspot and dollarspot; prior to the first snowfall to prevent snowmold. To control active diseases apply at first indication of damage. A second application may be required in one month if activity persists.

To prevent or control stripe smut and rust on bluegrass-fescue or ryegrass. Apply at **DOUBLE RATE**. Make applications in late fall and early spring to prevent stripe smut. To control active diseases apply at first signs of damage. A second application may be required in one month if activity persists.

In California use only on bluegrass turf and only in the period from late fall through early spring.

To prevent brownpatch on St Augustine, bahia and centipede grasses. Apply at **DOUBLE RATE** in early fall and in late winter or early spring. If disease activity persists repeat application in one month.

Not for use on predominately fine fescue turf.

Apply when foliage is dry. If temperatures are 80° or above, water lightly to remove granules from the grass blades.

Slight but temporary yellowing may occur following treatment but grass will recover with normal care.

ProTurf Division, O M Scott & Sons, Marysville, Ohio 43040