

NET WEIGHT 28½ LBS (12.75 kg)

8868



insecticide III



insecticide III



ProTurf

insecticide III  
turf insecticide

FOR PROFESSIONAL USE ONLY



insecticide III

ACCEPTED  
NOV 3 1977  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
this product is recommended for the pesticide  
category of turf insecticide.  
EPA Reg. No. 538-111



insecticide III  
turf insecticide

for use on turf/grasses or dichondra

**SUGGESTED SPREADER SETTINGS**  
to provide proper distribution utilize spreader before application

20% (11.75 lb) treats 22,000 sq ft (1/4 acre/2044 sq m) at NORMAL RATE  
25% (13.75 lb) treats 17,000 sq ft (1/3 acre/1527 sq m) at DOUBLE RATE  
28% (15.75 lb) treats 15,000 sq ft (1/3 acre/1372 sq m) at HEAVY RATE

SPREADER	SETTING OF TYPICAL	SPREADER SETTING	SPREADER SETTING NORMAL	SPREADER SETTING HEAVY
Becker (Crop Type)	3 mph	overlap wheels	24	24
Becker Rotary	3 mph	16 feet	14	14
Case-Lane and WPA	4.5 mph	20 feet	4.5	5.5
Bladen	4.5 mph	20 feet	4.5	5.5
Case-Lane	4.5 mph	20 feet	4.5	5.5
(PTO Model)	4.5 mph	20 feet	4.5	5.5
Case 8 ft A Series	4.5 mph	overlap wheels	15	17
Case 42" model	3 mph	overlap wheels	25	25

CONTROLS CHINCH BUG and other insects listed below:

- ant flea beetle (in dichondra)
- brown dog tick grasshopper
- cricket Hyperodes weevil
- cutworm sod webworm (lawn moth larvae)
- earwig vegetable weevil (in dichondra)

Apply over entire area. Overlap spreader wheel tracks so no strips are missed. Water as recommended after each application to wash product particles into the zone of insect activity. Do not mow or collect clippings for 24 hours after treatment.

CHINCH BUG control-apply at DOUBLE RATE any time activity or damage is noticed. Water thoroughly (1/2 to 1 inch) after application.

CHINCH BUG prevention-Apply at DOUBLE RATE as soon as daily temperatures reach 80° or above and repeat at 2 month intervals. Water thoroughly (1/2 to 1 inch) after each application.

SOD WEBWORM, CUTWORM, EARWIG, FLEA BEETLE and VEGETABLE WEEVIL control-apply at NORMAL RATE whenever insect activity is observed or suspected. Sprinkle lightly after application to remove product particles from the leaves.

ANT, TICK, CRICKET and GRASSHOPPER control-apply at DOUBLE RATE whenever insect activity is observed or suspected. Sprinkle lightly after each application to remove particles from leaves.

HYPERODES WEEVIL control-apply at HEAVY RATE as soon as grass greens in the spring and repeat in one month. Water lightly after application.

ProTurf Division, O M Scott & Sons, Marysville, Ohio 43040

8868

NET WEIGHT 28½ LBS (12.75 kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through the skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after using. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash all contaminated clothing with soap and hot water before reuse. Avoid contamination of food and foodstuffs and domestic water supplies. As directed, water lawn after application and allow to dry before permitting children or pets on treated areas. Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal. This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label. Do not reuse empty container. Destroy when empty.

Active ingredient: Chlorpyrifos O, O diethyl O (3, 5, 6-trichloro-2-pyridyl) phosphorothioate 1.34%  
Inert ingredients 98.66%  
Total 100.00%

EPA Reg No 538-111 EPA Est 538-OH-1 Product of USA  
ProTurf Division, O M Scott & Sons Marysville, Ohio 43040

8868



insecticide III

8868

538-111

11/3/1977

Page 131