

2



8774

NET WEIGHT 35 1/2 LBS (16.10 kg)

ProTurf

K-O-G weed control

8774

Scotts
ProTurf
K-O-G weed control

FOR PROFESSIONAL USE ONLY

NET WEIGHT 35 1/2 LBS (16.10 kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN
 Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Do not inhale dusts or vapors. Do not store near fire, stoves, radiators, furnaces, or other heat sources. Avoid application on garden plants, flowers and rootzone areas of trees and shrubs. Do not reuse container. Destroy by burning in a safe place away from water supplies. Do not burn.
 Active ingredient: Dicamba (2-Methyl-2,6-dichlorobenzoic acid) 93.0%
 Inert ingredients 6.0%
 Total 100.0%
 EPA Reg. No. 538-112 (EPA Est. 138-011)
 ProTurf Division, O M Scott & Sons, Marysville, Ohio 43040
 Product of USA
 Marysville, Ohio 43040

8774

ACCEPTED
SEP - 6 1979
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered Under EPA Reg. No. 538-112

8774



ProTurf

K-O-G weed control

8774

Scotts
ProTurf

K-O-G weed control

SUGGESTED SPREADER SETTINGS
 to provide proper distribution calibrate spreader before application

SPREADER	SPREADER RATE (lb/1000 sq ft)	SPREADER RATE (kg/ha)	SPREADER RATE (lb/1000 sq ft)	SPREADER RATE (kg/ha)
Scotts (Green Type)	3.0	6.7	2.0	4.5
Scotts (Blue Type)	3.0	6.7	2.0	4.5
Scotts (Red Type)	3.0	6.7	2.0	4.5
Scotts (Black Type)	3.0	6.7	2.0	4.5
Scotts (White Type)	3.0	6.7	2.0	4.5
Scotts (Yellow Type)	3.0	6.7	2.0	4.5
Scotts (Orange Type)	3.0	6.7	2.0	4.5
Scotts (Purple Type)	3.0	6.7	2.0	4.5
Scotts (Brown Type)	3.0	6.7	2.0	4.5
Scotts (Grey Type)	3.0	6.7	2.0	4.5
Scotts (Green Type)	3.0	6.7	2.0	4.5
Scotts (Blue Type)	3.0	6.7	2.0	4.5
Scotts (Red Type)	3.0	6.7	2.0	4.5
Scotts (Black Type)	3.0	6.7	2.0	4.5
Scotts (White Type)	3.0	6.7	2.0	4.5
Scotts (Yellow Type)	3.0	6.7	2.0	4.5
Scotts (Orange Type)	3.0	6.7	2.0	4.5
Scotts (Purple Type)	3.0	6.7	2.0	4.5
Scotts (Brown Type)	3.0	6.7	2.0	4.5
Scotts (Grey Type)	3.0	6.7	2.0	4.5

FOR BENTGRASS PUTTING GREENS
 For bentgrass putting greens apply at QUARTER RATE to moist foliage for control of clover and chickweed. Repeat treatment in 3 weeks if needed. (Treats 44,000 sq ft)

FOR USE ON OTHER TURFGRASS AREAS
 Turfgrass areas other than greens apply to moist or dry foliage at—

HALF RATE to control clover (treats 22,000 sq ft).
NORMAL RATE to control knotted, chickweed (common and mouse-ear), sheep sorrel and black medic (treats 11,000 sq ft).

DOUBLE RATE to control wild onion and garlic, curly dock, ground ivy, Canada thistle, cudweed, yarrow and horseweed. For control of yarrow and wild onion and garlic start treatments in either spring or fall and repeat at 6 month intervals for a total of 2 or 3 applications as needed (treats 5,500 sq ft).

Avoid application on rootzone areas of trees and shrubs. Apply anytime weeds are actively growing. Do not apply when ground is frozen.

Delay planting grass seed until after a heavy rain or watering and not sooner than two weeks after treatment.

On new plantings delay treatment until after the second mowing.

Not for use on St. Augustine grass or Dichondra.
 ProTurf Division, O M Scott & Sons, Marysville, Ohio 43040

8774