

538-112

PM 25

1061

8774

K-O-G
Weed Control

8774

K-O-G
Weed ControlK-O-G
Weed Control

DO NOT USE

EPA Form 8570-13 (3-75)

K-O-G
Weed ControlK-O-G
Weed
Control

ACCEPTED

FEB 4 1982

Under the Federal Insecticide
Fungicide and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 538-112K-O-G
Weed Control

•provides postemergence weed control

Net Weight 35 1/2 lbs (16.10 kg)

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENT

Hazardous to Fish and Aquatic Life. May Cause
Chronic Effects. Avoid contact with plants, animals,
and children. Do not use in residential areas.
Do not use on lawns, lawns, or other areas where
children or pets play.

EPA REG. NO. 538-112

Aerial Application

Application Rate

Application Rate

Application Rate

Application Rate

Application Rate

Application Rate

Application Rate

8774

For Professional Use Only

8774

8774

K-O-G
Weed Control

8774

K-O-G
Weed Control

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Delay planting grass seed until after a heavy rain or watering and not sooner than two weeks after treatment. Apply any time weeds are actively growing. Do not apply when ground is frozen.

For bentgrass putting greens

For bentgrass putting greens apply at QUARTER RATE to most foliage for control of clover and chickweed. Repeat treatment in 3 weeks if needed. (Treats 44,000 ft².)

For use on other turfgrass areas

Turfgrass areas other than greens apply to moist or dry foliage at—

HALF RATE to control clover (treats 22,000 ft².)

NORMAL RATE to control knotweed, chickweed (common and mouse-ear), sheep sorrel and black medic (treats 11,000 ft².)

DOUBLE RATE to control wild onion and garlic, curly dock, ground ivy, Canada thistle, cutweed, yarrow and horseweed. For control of yarrow and wild onion and garlic start treatments in either spring or fall and repeat at 6 month intervals for a total of 2 or 3 applications as needed (treats 5,500 ft².)

Do not apply when ground is frozen. Delay planting grass seed until after a heavy rain or watering and not sooner than two weeks after treatment.

On new plantings delay treatment until after the second mowing. Not for use on St. Augustinegrass or Dichondra.

Avoid application to garden plants, flowers, and rootzone areas.

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Pesticide opened bags by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a secondary landfill or by incineration if allowed by State and local authorities.

Note

Avoid application on rootzone areas of trees and shrubs.

Proflurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

35 1/2 lb (16.10 kg) treats 44,000 ft² (4 acre) at QUARTER RATE

35 1/2 lb (16.10 kg) treats 22,000 ft² (2 acre) at HALF RATE

35 1/2 lb (16.10 kg) treats 11,000 ft² (1 acre) at NORMAL RATE

35 1/2 lb (16.10 kg) treats 5,500 ft² (1/2 acre) at DOUBLE RATE

| SPREADER | CALIBRATION | SPREADER | SPREADER SETTINGS | | | |
|--------------------|-------------|----------|-------------------|-------|-------|-------|
| | | | Rate | Rate | Rate | Rate |
| Scotts (Drop Type) | 3 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |
| Scotts Rotary | 3 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |
| Leifherr | 4.5 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |
| W/R Skyles | 4.5 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |
| Leifherr | 4.5 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |
| P/D Skyles | 4.5 mph | 16 feet | 3 1/2 | 4 1/2 | 5 1/2 | 7 1/2 |