

8774

K-O-G
Weed Control

8774

K-O-G
Weed ControlK-O-G
Weed Control

Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply only those weeds are actively growing. Do not apply when ground is frozen.

For bentgrass putting greens

For bentgrass putting greens apply at QUARTER RATE to avoid damage for control of annual and perennial. Repeat treatment in 3 weeks if needed. (Rate 0.500 lb/1000 sq ft)

For use on other turfgrass areas

Turfgrass areas other than greens apply to moist or dry foliage at—

HALF RATE to control clover (Rate 2.000 lb/1000 sq ft)

NORMAL RATE to control broadleaf, annual broadleaf and spurge-cacti, along control and thick mats (Rate 4.000 lb/1000 sq ft)

DOUBLE RATE to control wild onion and garlic, early dock, ground ivy, Canada thistle, nutgrass, yellow and hairgrass. For control of yellow and wild onion and garlic start treatment in either spring or fall and repeat at 8 month intervals for a total of 2 or 3 applications as needed (Rate 8.000 lb/1000 sq ft)

Do not apply when ground is frozen.

Delay planting grass seed until after a heavy rain or watering and not earlier than two weeks after treatment.

On new plantings delay treatment until after the second mowing.

Not for use on St. Augustinegrass or Stenotaphrum.

Avoid application to garden plants, flowers, and vegetable areas.

Storage and disposal

STORAGE: Store in a clean, dry place. Re-seal opened bag by folding top-down and securing.

POISONED DISPOSAL: Do not contribute water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Note

Avoid application on sensitive areas of lawns and shrubs.

Proflurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

WEED TYPE	Suggested Spreader Settings			
	SPREADER TYPE	SPREADER RATE (lb/1000 sq ft)	SPREADER RATE (lb/1000 sq ft)	SPREADER RATE (lb/1000 sq ft)
Annual Broadleaf	1/2" - 3/4"	2.00	4.00	8.00
Wild Onion and Garlic	1/2" - 3/4"	4.00	8.00	16.00
St. Augustinegrass	1/2" - 3/4"	2.00	4.00	8.00
Stenotaphrum	1/2" - 3/4"	2.00	4.00	8.00

K-O-G
Weed ControlK-O-G
Weed
ControlK-O-G
Weed Control

-provides postemergence weed control

Net Weight 35 1/2 lbs (16.10 kg)

KEEP OUT OF REACH OF CHILDREN
WARNING

PRECAUTIONARY STATEMENT

Hazardous to Humans and Domestic Animals (antidotes available). Contains substances that are toxic to fish. Do not use in or on streams, rivers, ponds, lakes, or other bodies of water. Do not use in or on areas where the product may come into contact with water. Do not use in or on areas where the product may come into contact with water. Do not use in or on areas where the product may come into contact with water.

If you experience any adverse effects, stop using the product immediately and consult your physician. Do not use in or on areas where the product may come into contact with water. Do not use in or on areas where the product may come into contact with water.

Environmental Precautions: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

ACTIVE INGREDIENTS

Quantity of Active Ingredient 0.2%

Other Ingredients 99.8%

Net Weight 35 1/2 lbs (16.10 kg)

Net Content 35 1/2 lbs (16.10 kg)

Product Group: The O M Scott & Sons Company
Marysville, Ohio 43041

8774

8774

For Professional Use Only

8774

K-O-G
Weed Control

NOT REVIEWED
In Accordance with PR Notice 82-
Based on Draft Labeling Dated 8/19/87

8774

K-O-G
Weed Control

u/1/88