

10 days
100%
100%



101V broad spectrum fungicide

Controls these turfgrass and dichondra (in California only) diseases:

- leafspot (Bipolaris)
- dollar spot
- brown patch
- copper spot
- curvularia leafspot
- rust (stem rust on bluegrass)
- alternaria leafspot (in Dichondra)
- gleosporium (anthracnose in Dichondra)

USE THESE SPREADER SETTINGS to ensure proper distribution

29 Lbs (13.2 Kg) treats 11,000 sq ft (1 Acre/1022 sq m) at NORMAL RATE
29 Lbs (13.2 Kg) treats 5,500 sq ft (1/2 Acre/511 sq m) at DOUBLE RATE

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	6'	9'
Gandy 8 R A Series	4.5 mph	overlap wheels	25	34
Gandy 42 Model	2.5 mph	overlap wheels	35	44

FOR USE ON TURFGRASS OR DICHONDRA

To prevent diseases apply to moist foliage at NORMAL RATE at 10-14 day intervals. Begin applications at expected time of disease activity and continue as climatic conditions remain favorable for disease development.

To control active diseases apply to moist foliage at NORMAL RATE at first signs of disease. Follow the preventative application schedule to bring the disease under control and prevent further infestation.

For severe disease infestations make first application to moist foliage at DOUBLE RATE and follow at 7-10 day intervals at NORMAL RATE until disease activity is arrested.

PRECAUTION

Please pay special attention to the handling information as noted in the caution statement to ensure safe use of the product.

ProTurf Division, OMScott & Sons Co, Marysville, Ohio 43040