

G Weed control 8774

ACCEPTED
AUG 29 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

Lely-HR	450 rpm	10 feet				
(PTO Model)	@4.5 mph					
Gandy 8 ft A Series	4.5 mph	overlap wheels	12	15½	20	26
Gandy 42" Model	2.5 mph	overlap wheels	24	30	37	45

FOR BENTGRASS PUTTING GREENS

For bentgrass putting greens apply at **QUARTER RATE** to moist foliage for control of clover and chickweed. Repeat treatment in 3 weeks if needed. (Treats 44,000 sq ft)

FOR USE ON OTHER TURFGRASS AREAS

Turfgrass areas other than greens apply to moist or dry foliage at—

HALF RATE to control clover (treats 22,000 sq ft).

NORMAL RATE to control knotweed, chickweed (common and mouse-ear), sheep sorrel and black medic (treats 11,000 sq ft).

DOUBLE RATE to control wild onion and garlic, curly dock, ground ivy, Canada thistle, cudweed, yarrow and horseweed. For control of yarrow and wild onion and garlic start treatments in either spring or fall and repeat at 6 month intervals for a total of 2 or 3 applications as needed (treats 5,500 sq ft).

Avoid application on rootzone areas of trees and shrubs.

Apply anytime weeds are actively growing. Do not apply when ground is frozen.

Delay planting grass seed until after a heavy rain or watering and not sooner than two weeks after treatment.

On new plantings delay treatment until after the second mowing.

Not for use on St Augustinegrass or Dichondra.

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040

8774


Handwritten notes and a stamp in the top right corner, including the number 53.

XXXX A JoluoC
annua eod

XXXX A JoluoC
annua eod

XXXX A JoluoC
annua eod


NET WEIGHT 24¾ LBS (11.2 Kg)



poa annua cont



**poa annua
control V**
controls poa annua in bermudagrass turf



poa annua cont

poa annua control V

XXXX

poa annua control V

controls poa annua in bermudagrass turf

FOR PROFESSIONAL USE ONLY

NET WEIGHT 24³/₄ LBS (11.2 Kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid skin and eye contact. In case of skin contact wash with soap and water. In case of eye contact, wash with large quantities of water.

Apply only as specified on this label. Do not reuse empty container. Destroy when empty.

Active ingredient:		0.90%
3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide		99.10%
Inert ingredients:		Total 100.00%

EPA Reg No 538-1-5	EPA Est No 538 OH 1	Product of USA
ProTurf Division, C. M. Scott & Sons		Marysville, Ohio 43040

poa annua control V

XXXX

XXXX


Feb. 21, 1975

ORDER # 538-115

XXXX poa annua control V

XXXX poa annua control V

TO OPEN BAG
REMOVE TOP



poa annua control V



poa annua control V

Controls poa annua in bermudagrass turf.
Also controls bluegrass, bentgrass and other cool season grasses.

USE THESE SPREADER SETTINGS to ensure proper distribution

24 3/4 Lbs (11.2 Kg) treats 22,000 sq ft (1/2 Acre/2045 sq m) at NORMAL RATE
24 3/4 Lbs (11.2 Kg) treats 11,000 sq ft (1/4 Acre/1022 sq m) at DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	5	7
Gandy 8 ft A Series	4.5 mph	overlap wheels	18	25
Gandy 42" Model	2.5 mph	overlap wheels	29	36

For use on bermudagrass only.
Apply to moist or dry foliage. Water thoroughly after application to move the active ingredient into the rootzone.
For control of Poa annua, apply at NORMAL RATE any time it is actively growing after germination in the fall thru the spring.
For elimination of winter overseeded cool season grasses and

Poa annua control V

XXXXX

**USE THESE SPREADER SETTINGS
to ensure proper distribution**

24 3/4 Lbs (11.2 Kg) treats 22,000 sq ft (1/2 Acre/2045 sq m) at NORMAL RATE
24 3/4 Lbs (11.2 Kg) treats 11,000 sq ft (1/4 Acre/1022 sq m) at DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	5	7
Gandy 8 ft A Series	4.5 mph	overlap wheels	18	25
Gandy 42" Model	2.5 mph	overlap wheels	29	36

For use on bermudagrass only.

Apply to moist or dry foliage. Water thoroughly after application to move the active ingredient into the rootzone.

For control of Poa annua, apply at NORMAL RATE any time it is actively growing after germination in the fall thru the spring.

For elimination of winter overseeded cool season grasses and conversion to bermudagrass, on golf course tees and greens, apply at NORMAL RATE in the spring when weather favors bermudagrass regrowth.

For control of Kentucky bluegrass and bentgrass, in areas other than tees and greens, apply at DOUBLE RATE any time they are actively growing.

PRECAUTIONS

Do not use on dichondra, bluegrass, annual or perennial ryegrass, fescuegrasses, bentgrass or bahiagrass where these grasses are the desired turf.

Not for use on muck or peat soils.

Do not apply this product where winter overseeding with cool season turfgrasses such as bluegrasses, ryegrasses, bentgrasses or fescuegrasses is planned within 90 days.

Avoid treating areas on fairways or approaches that drain into bentgrass greens or to areas that have been overseeded with cool season grasses.

Application on golf greens prior to a severe cold or freezing period can cause discoloration. Avoid winter applications where these conditions are likely to occur.

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040

XXXXX