



19

XXXX

Treats 2750 sq ft

Scotts

ProTurf

NEW 25-0-0
Super Bonus
for dichondra

weed and insect control plus fertilizer for dichondra

FOR PROFESSIONAL USE ONLY IN CALIFORNIA

DO NOT USE ON GRASS LAWNS

INSTRUCTIONS FOR USE

Two applications (6 to 8 weeks apart) any time of the year provides control of the following weeds when they are actively growing plus prolonged feeding. Also controls insects shown when active (spring/summer).

Grassy Weeds	Broad Weeds	Insects
bentgrass bluegrass (suppression only) coarse fescue (suppression only)	foxtail barley Poa annua redtop ryegrass velvetgrass	chickweed common mousetar cudweed hairy
		asails pearwort saliva sedges white clover
		sod webworm beetles cutworms

WINTER SPRING PROGRAM—controls above weeds plus—
• prevents spotted spurge and crabgrass
• delays spring revival of bermudagrass
• controls vegetable weevil larvae

Apply in December, January or early February and repeat in 6 to 8 weeks. To prevent crabgrass and spurge both applications must be made before mid March in Southern California, late March in the Interior Valley and before mid April in Northern California.

FALL PROGRAM—to prevent Australian brass buttons, chickweed, foxtail barley, mallow, Poa annua and shepherd's purse make an additional single application in September, October or early November.

To ensure proper distribution
use a Scotts Spreader of setting # 7

Slightly overlap spreader wheel tracks to provide complete coverage.

PRECAUTIONS

Apply only to dry foliage. Important to rinse particles off leaves by adequate irrigation or hand hose spray immediately after application.

A new dichondra seeding can be treated with Super Bonus when the dichondra has covered the soil but no sooner than 2 months after seeding. Wait 4 weeks after sodding or plugging. After application delay planting of any dichondra for 8 weeks.

Avoid application on shaded area receiving less than 4 hours of sunlight daily.

Avoid application on shrubs, flowers, or vegetable plants.

NET WEIGHT 17½ LBS (8.0 Kg)

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. May be absorbed through skin. Avoid inhalation in contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after use. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash all contaminated clothing with soap and hot water before reuse. Avoid contamination of food and foodstuffs. Avoid treated water lawn after application and allow to dry before permitting children or pets on treated areas. Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by washing of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste.

Active ingredients

Diflufenican (1,1-Dimethyl-2,2-dicyclopentyl-3,3-difluoropropane)	3.5%
Metolachlor (1-Butyl-3-(3,4-dichlorophenyl)-1-methylurea)	1.06%
Miflumuron (3-(p-chlorophenyl)-1,1-dimethylurea)	0.17%
Chlorpyrifos (0,0-diethyl 0,1,3,5,6-trichloro-2-pyridyl phosphorothioate)	0.26%
Inert ingredients	95.01%
	Total 100.00%

EPA Reg No 538-121-AA EPA Est 538-CH-1
Net weight 17½ lbs ProTurf New Super Bonus for Dichondra
25.00 guaranteed analysis

Total nitrogen (N) 25.00%
16.0% water-soluble nitrogen from urea and methylene ureas
8.4% water-soluble nitrogen from methyl ureas
Potential acidity: 600 lbs/acre lim carbon (lime per ton)
US Process Pat No 3,076,700 US Composition Pat No 3,083,089 Product of USA
ProTurf Division, O.M. Scott & Sons Company, Marysville, Ohio 43040

June, 3, 1975

538-121

BEST DOCUMENT AVAILABLE

XXXX

53