

as amended, for the pesticide registered under EPA Reg. No. 261-10-1

XXXX

XXXX

XXXX

NET WEIGHT 41½ LBS (18.8 Kg)

XXXX

XXXX

TO OPEN BAG

ProTurf
101 broad spectrum fungicide

FOR PROFESSIONAL USE ONLY

ProTurf
101 broad spectrum fungicide

XXXXX

NET WEIGHT 41½ LBS (18.8 Kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Can be irritating to eyes or skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and warm water after handling. In case of contact with eyes, flush with plenty of water for at least 15 minutes. Do not store near feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Destroy by burying or burning in a safe place. Stay away from smoke or fumes.

Active ingredient Chlorothalonil (Tetrachloroisophthalonitrile)	7.40%
Inert ingredients	92.60%
Total 100.00%	

EPA Reg. No. 538-124-A-A EPA Est. 53R-081-1 Product of USA
ProTurf Division, O.M. Scott & Sons Co. Marysville, Ohio 43040

XXXXX

ProTurf
101 broad spectrum fungicide

Controls these turfgrass diseases:
leafspot (Holminthoerium) copper spot
dollarspot stem rust of bluegrass
brownpatch curvularia leafspot

Controls these dichondra diseases (in California only):
alternaria leafspot
gleosporium (anthracnose)

SUGGESTED SPREADER SETTINGS
To ensure proper distribution calibrate spreader before application.
41½ lbs (18.8 Kg) treats 11,000 sq ft (¼ Acre—1022 sq m)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS
Scotts (Drop Type)	3 mph	overlap wheels	6 ¼"
Scotts Rotary	3 mph	8 feet	1"
Lely—WTR & WFR Models	4.5 mph	16 feet	5 ½" X
Lely—HR (PTO Model)	4.5 mph	16 feet	5 ½" X
Gandy B.P.A. Series	4.5 mph	overlap wheels	22"
42" Model	2.5 mph	overlap wheels	46"

FOR USE ON GOLF COURSE TEES, GREENS AND DICHONDRA

To prevent diseases apply to moist foliage at 10-14 day intervals. Begin applications at expected time of disease activity and continue as climatic conditions remain favorable for disease development.

To control active diseases apply to moist foliage at first signs of disease. Follow the preventative application schedule to bring the disease under control and prevent further infestation.

For severe disease infestations apply to moist foliage every 7-10 days until the disease is stopped.

PRECAUTION

Please pay special attention to the handling information as noted in the caution statement to ensure safe use of the product.

ProTurf Division, O.M. Scott & Sons Co. Marysville, Ohio 43040