

XXXX

Treats 2750 sq ft



Bonus for dichondra

Weed control plus fertilizer for dichondra

HOW TO USE

To control these actively growing weeds, apply any time and repeat in 8 weeks:

oxalis	filaree	soloiva
white clover	cudweed	poa annua
chickweeds	pearlwort	annual plantain

To prevent spotted spurge, apply in December, January or early February and repeat in 8 weeks.

To prevent crabgrass, apply in December, January, February or March and repeat in 8 weeks.

To ensure proper distribution
use a Scott Spreader at setting #7

Slightly overlap spreader wheel tracks to provide complete coverage.

PRECAUTIONS

Completely rinse particles off leaves immediately after application by adequate irrigation or hand hose spray.

Do not use on grass lawns.

On newly seeded areas wait 6 months before treating, on newly sodded or plugged areas wait one month. After application, wait 6 months before reseeding.

Do not use on shaded areas receiving less than half a day of sunlight.

This product does not control deep-rooted perennials such as bermuda-grass, paspalums, dandelions and coarse fescue.

Net Weight 12½ Lbs (5.7 Kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN
Avoid contact with skin, eyes and clothing. May cause irritation of eyes, skin, nose or throat. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

Active ingredient: Neburon 1-Butyl-3-(3,4-dichlorophenyl)-

1-methylurea	2.00%
Inert ingredients:	98.00%
	Total 100.00%

EPA Reg No 538-125-AA EPA Est 538 OH-1

Net weight 12½ lbs ProTurf Bonus for Dichondra 35-0-0 guaranteed analysis

Total nitrogen (N) 35.00%

23.30% water soluble nitrogen from urea and methylene ureas

11.70% water insoluble nitrogen from methylene ureas

Available phosphoric acid (P.O.) 0.00%

Soluble potash (K.O.) 0.00%

Potential acidity equivalent to 1350 lbs calcium carbonate per ton


US Process Pat Nos 3,231,363; 3,705,794 Other Pat Pend Product of USA

ProTurf Division, O M Scott & Sons Co Marysville, Ohio 43040

XXXX

4
12/12/74
L-25-1457
TEST DOCUMENT AVAILABLE

• Con
• Prov
• Lepre
• Catfish
• South
• Tuna
• Nettles
DO App
Dec 01



3436

ntact with
ster using.
i. Wash all
and food-
or pets on
antidotal.

it contam-
is likely to

3.50%
0.12%
1.06%
0.26%
95.01%
100.00%
ed analysis
25.00%

Ohio 43040

3436



A triangular logo with the words "EASY OPEN BAG" at the top and "WHITE TAPE" at the bottom.

6052

Treats 5,000 sq ft

The logo features the Scotts brand name in its signature script font inside a dark oval shape. To the right of the oval, the word "Kansel" is written in a large, bold, serif font. Below "Kansel", the words "lawn weed control" are written in a smaller, sans-serif font.

controls dandelions and these lawn weeds

How To List

- DON'T** use when weeds are up and growing (serve as the bait). Lawns may wait until after the second mowing of the year but no later than late March. In the fall, don't treat after the first of November.

DON'T apply only when lawn is wet from new rain or thorough watering. Very important so that the Scott's Kassel particles will stick to the weed leaves instead of falling to the soil.

DON'T withhold watering for at least 24 hours after application to wash particles off the weed leaves.

DON'T apply on a calm day.

DON'T apply if Scott's Spreader (Standard or Automatic) Model setting treats 1,000 sq ft. **5**

On bentgrass and permu-
tgrass lawns, use setting **3½**

DON'T apply when rain is expected within 24 hours.

DON'T use when grass is wet or touches bottom of spreader. Mow lawn first in such weather.

DON'T use on a newly seeded, sodded or sprouted lawn. Wait until it has been mowed at least 4 times.

DON'T apply just after using Kassel. Wait at least two weeks and after applying a rain.

DON'T use on St. Augustine grass, carpet grass, dichondra, lawn or tropical and flowers, shrubs, fruit trees, trees, etc.

DON'T apply on the coarse areas of shrubs or under the branch spread of trees.



卷之三



6052

Kamse!

lawn weed control



6052

Kyansel

6052

SCOTT'S NO-(U)IBBLE (SUBSTITUTE)

bulletin-service

that to expect

6052




Treats 5,000 sq ft




Scotts
Kansel
lawn weed control

controls dandelions,
buckhorn,
chickweed, knotweed,
plantain, white clover



Kansel
lawn weed control



Kansel
lawn weed control

controls dandelions,
buckhorn,
chickweed, knotweed,
plantain, white clover
and other weeds

(see complete list on the back panel)

Not for use on St Augustinegrass,
carpetgrass or dichondra lawns

A problem-solving product from Scotts, the lawn people

NET WEIGHT 5 LBS (treats 5,000 sq ft)

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. In case of contact flush with water. For eyes get medical attention. Avoid application near or over garden plants and flower. Keep away from rootzone areas of shrubs and the ground spread of trees. Do not reuse container. Dispose when empty.

Do not contaminate water used for irrigation or domestic purposes.

Active ingredients: 24-Dichlorophenoxyacetic acid	2.95
Dicamba (Monsanto 3,6-dichlorobenzoic acid)	1.45
Inert ingredients	96.57
EPA Reg. No. 5000-CAV (EPA Est. #3-001)	Total 100.00
US Composition Part No. 3003000	Product of USA
OMRI Listed & Safe	Maryland Origin 100%

6052

6052

6052



Kansel
lawn weed control

Kansel
lawn weed co



XXXX

Treats 2750 sq ft



35-0-0

Bonus for dichondra

Weed control plus fertilizer for dichondra

HOW TO USE

To control these actively growing weeds, apply any time and repeat in 6 weeks:
grass, white clover, chickweed, purple annua, vetch, chickweed, peachroot, prunus plantain
To prevent applied spur 2A, apply in December, January or early February and repeat in 6 weeks.
To prevent crabgrass, apply in December, January, February or March and repeat in 6 weeks.

To ensure proper distribution
use a Scott spreader at setting 7

Slightly overlap spreader wheel tracks to provide complete coverage

PRECAUTIONS

Completely rinse particles off leaves immediately after application by adequate irrigation or hand hose spray.
Do not use on grass lawns
On newly sown areas wait 6 months before treating, on newly sodded or plugged areas wait one month. After application, wait 6 months before reseeding.
Do not use on shaded areas receiving less than half a day of sunlight.
This product does not control deep-rooted perennials such as Bermuda grass, periwinkles, dianthus and coarse fescue.

Net Weight 12½ Lbs (5.7 Kg)

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes and clothing. May cause irritation of eyes, skin, nose or throat. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

Active ingredient: Neburon 1 Butyl-3-(3,4-dihydrophenyl)-

1-methylnaphthalene, 2.00%

Inert ingredients, 98.00%

Total 100.00%

EPA Reg No 538-125 AA EPA Est 538 OH 1

Net weight 12½ lbs ProTurf Bonus for Dichondra 35.00 guaranteed analysis

Total nitrogen (N) 35.00%

23.30% water soluble nitrogen from urea and methylene ureas

11.70% water insoluble nitrogen from methylene ureas

Ammonium phosphate (P.O.) 0.00%

Soluble potash (K O) 0.00%

Potential acidity equivalent to 1350 lbs calcium carbonate per ton

US Process Pat Nos 3,231,363, 3,705,794 Other Pat Pend Product of USA

ProTurf Division O M Scott & Sons Co Marysville, Ohio 43040

XXXX