

JUN 30 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 538-128

Vegetable Garden Weed Preventer

COVERS 1500 SQ. FT. SANDY SOIL
1100 SQ. FT. HEAVY SOIL

Prevents many undesirable weeds and grasses in vegetable gardens including:

- | | | | |
|------------------|------------------|--------------------------|-----------------|
| barnyardgrass | crabgrass smooth | groundcherry | burstlane |
| browntop panicum | dodder | johnsongrass (from seed) | redroot pigweed |
| carpetweed | Florida pusley | rambouniers | sandbur |
| chickweed | goosegrass | lovegrass | witchgrass |
| crabgrass large | green foxtail | poa annua | yellow foxtail |

NEW!



Scott's Vegetable Garden Weed Preventer

Scott's Vegetable Garden Weed Preventer

Prevents many undesirable weeds and grasses in vegetable gardens (see back)

Scott's Vegetable Garden Weed Preventer

ADVANTAGES OF SCOTT'S® VEGETABLE GARDEN WEED PREVENTER:

Minimizes early season weed competition. Prevents many undesirable weeds and grasses in vegetable gardens (see list)

Will not injure crops listed when used as directed. Leaves no harmful residue when used as directed. No-Quibble Guarantee: If for any reason you are not satisfied, send evidence of purchase and Scott's will refund your money.

EASY TO USE!

When: Apply at time of seeding or transplanting:

- beans (dry and snap)
- black-eyed (southern) peas
- broccoli
- brussels sprouts
- cabbage
- cauliflower
- chinese cabbage
- collards
- garlic
- kale
- mustard greens
- onions
- potatoes (except California)
- strawberries (before first bloom)
- sweet corn
- sweet potatoes
- turnips
- yams



Apply when plants are well established.

- cucumbers, melons, squash (after plants have 4-5 true leaves)
- eggplants, peppers, tomatoes (4-6 weeks after transplanting or when seeded plants are 4-6 inches tall)



Where: in seed or transplant bed or around established plants

How:

- Cultivate soil and eliminate existing weeds
- Fertilize the soil, then seed
- Apply this product uniformly
- Do not rake in
- Water lightly after application



How Much:

- On sandy soils (all listed crops except strawberries): 1 cup (0.25 lb) on 75 sq ft Scott's spreader setting 7
- On heavy soils (all listed crops except strawberries): 1 cup (0.25 lb) on 55 sq ft Scott's spreader setting 8 1/2
- On any soil (strawberries only): 1 cup (0.25 lb) on 65 sq ft Scott's spreader setting 7 1/2

Application Rate (Actual-Size)

Sprinkle evenly as shown



Sandy Soils

Heavy Soils

Replanting Notes:

- All crops listed above may be planted following harvest of a treated crop
- Replanting crops other than those listed above in treated soil within 8 months of application may result in crop injury
- If replanting is required because of early crop failure, the planting of listed crops may result in crop injury

Precautions:

- Wash vegetables before eating
- Do not use on beets, lettuce, cow peas, Tennessee greenpod, snap beans and other crops not listed on this label
- Does not control established weeds
- Disturbing soil after application reduces effectiveness



SCOTT'S® NO-QUIBBLE GUARANTEE

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

O.M. Scott & Sons • Marysville, Ohio 43040



CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Avoid contact with eyes, skin and clothing. In case of contact, flush eyes with plenty of water. Wash after handling or using. Do not graze treated areas or feed plant parts to livestock. Do not contaminate feed, foodstuffs, seed and fertilizers or water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of

wastes. Apply this product only as specified on this label. Do not reuse container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Active ingredient: Dimethyl trichloroterephthalate 5.00%
Inert ingredients: 95.00%
Total: 100.00%

NET WT 5 lbs

EPA Reg No. 538-128
EPA Est. 34637-041



- Use:
- Vegetable Garden
 - Vegetable Garden
 - Vegetable Garden

DIRECTIONS FOR USE

HOW TO USE: Apply at the rate indicated below. Spray when conditions for growth of turf grasses are good. . . During droughts, water several days before treating and fertilize to restore green color. Keep lawn mowed 1½ to 2 inches during treatment. Spray newly seeded lawns at "Seedling" rate shown below after they are well established. Do not use on St. Augustine, Bahia Carpet or Centipede grass.

CAN BE USED ON:

TYPE OF LAWN	PESTS	DIRECTIONS
New lawns, more sensitive grasses, such as Fescues and Bentgrass in early Spring or late Fall during cool days and moist soil conditions.	Crabgrass in "seedling" stage, Witchgrass, Barnyardgrass.	Spray at rate of 14 tbsps. or 7 ounces to each 5 gallons of water per each 1,000 sq. ft. (25 ft. x 40 ft.) of lawn in early Summer when crabgrass is young. Apply 2 or more treatments at 7 day intervals. If new crabgrass growth appears later in the season, repeat applications may be necessary.
	Crabgrass in "branching" stage.	Spray at rate of 14 tbsps. to 5 gallons of water per each 700 sq. ft. (20 ft. x 35 ft.) Apply 2 or more treatments at 7 day intervals.
Established lawns, such as Kentucky, Merion Blue, Dichondra, Zoysia, Bermuda.	Crabgrass in "branched" stage, Witchgrass, Fox tail, Barnyardgrass, Dallis grass, Sandbur, Chickweed.	Spray at rate of 24 tbsps. or 12 ounces to 5 gallons of water per each 1,000 sq. ft. (25 ft. x 40 ft.). Repeat spraying in 7 days. Application may discolor, but will not permanently injure desirable grasses.

On non-crop or non-pasture areas, such as drainage ditch banks, fence rows, rights of way, to control Johnson grass, dallis grass, sandburs, barnyardgrass, spray on grass to point of run off diluted with water at the rate of 1 pint in 4 gallons of water, 1 quart in 8 gallons of water and 1 gallon in 32 gallons of water.

CAUTION: Harmful if swallowed. If taken internally, induce vomiting. CALL A PHYSICIAN AT ONCE. Avoid contact with skin or breathing of spray mist. Keep away from children and domestic animals. Avoid contamination of feed and foodstuffs. Wash hands thoroughly after using. Do not contaminate water used for irrigation or domestic purposes.

Keep livestock off treated area.

Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.

11507

U.S.D.A. Reg. No. 539-100

DISTRIBUTED BY SEARS, ROEBUCK AND CO.
Chicago, Illinois

**LIQUID
CRABGRASS
KILLER**

ALSO KILLS: FOXTAIL, CHICKWEED, DALLIS GRASS, BARNYARDGRASS, SANDBUR, WITCHGRASS, JOHNSON GRASS

DO NOT USE ON LAWNS OTHER THAN St. Augustine, Centipede, Carpet and Bahia grass and NON-CROP AREAS.

ACTIVE INGREDIENTS:

Disodium Methanearsonate	12.6%
INERT INGREDIENTS:	87.4%
TOTAL	100.0%

Total Arsenic as elemental, all in water soluble form 5.13%.
Equivalent to 20.0% Disodium Methanearsonate Hexahydrate.

NET CONTENTS 1 PINT

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
FOR ADDITIONAL CAUTIONS SEE BACK PANEL.**

ACCEPTED
May 11, 1972
APPROVED BY FEDERAL INSPECTOR
FOR ECONOMIC ACTION
ED UNDER NO. 539-100

