

FRONT:

1989 Package Copy for #XXXX
VEGETABLE GARDEN WEED PREVENTER
05/04/88 AR 05/05/88 AR
05/10/88 AR 06/06/88 AR
08/26/88 AR 08/23/88 AR/VS
05/23/89 VS 04/07/92 VS/cr

(Photo)

(Scotts (R))

Vegetable Garden
Weed Preventer

Weeds 665 sq ft
Example - 5 ft. by 133 ft.

Sprinkle around your Flowers, Roses, Shrubs
Trees, Groundcovers and Vegetables and prevent weeds
this summer

*Contains Dacthal

Net. Wt. 2.75 lbs.

ACTIVE INGREDIENTS:

DCPA (dimethyl tetrachloroterephthalate)	By Wt.
	5.0%
INERT INGREDIENTS:	95.0%
	Total 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back for additional precautionary statements

*Dacthal is a registered trademark of Fermenta Plant Protection Company

EPA Reg. No. 538-128


EPA Est XXX-XX-X

Product of USA

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1990, The O M Scott & Sons Company. All rights reserved.



Scotts Vegetable Garden Weed Preventer is a pre-emergence application for control of crabgrass, foxtail, chickweed and certain other annual weeds in Flowers, Roses, Shrubs, Trees, Groundcovers, Strawberries and Vegetables. See folder on side of container for directions and complete list of plants where this product may be used.

(Photo)

(Treated)

(Untreated)

(Photo) Sprinkle the granules over the soil surface at the rate shown in this photo which equals 1 cup per 60 sq ft (6' x 10'). ✓

**Scotts Policy
No-Quibble Guarantee**

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.
Scotts...the lawn & garden people
Marysville, Ohio 43041

0 (UPC)

32247 02925

Questions about weeds in your garden? Call Scotts Toll-Free HOTLINE:
1-800-543-TURF (In Ohio 1-800-762-4010)

Directions For Use

Carefully read and follow instructions inside this folder

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans And Domestic Animals. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. If in eyes: Flush with plenty of water. Call a physician if irritation persists. Do not graze treated areas or feed plant parts to livestock.

ENVIRONMENTAL HAZARDS

Collect and incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes.

Groundwater Advisory: Tetrachloroterephthalic acid, a breakdown product of Dacthal, is known to leach through soil as a result of agricultural turf uses and has been found in groundwater which may be used for drinking water. Users are advised not to apply Dacthal to sand or loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater used for drinking water.

STORAGE: Store in a clean, dry place. Reseal opened container.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse packaging before discarding in trash.

Vegetables:

Broccoli, Brussels Sprouts, Cauliflower, Cabbage, Garlic, Onions, Sweet Potatoes: Apply at time of seeding or transplanting.

Beans (dry or snap), Southern Peas (blackeyed), Collards, Kale, Mustard Greens, Turnips: Apply at time of seeding.

Seeded Cucumbers, Squash (winter and summer), Melons (cantaloupe, watermelon, honeydew): Apply after plants are well established and have 4 to 5 true leaves, and when growing conditions are favorable for good plant growth. Weed prior to application if weeds have emerged.

Potatoes (whole or cut pieces): Apply at time of planting. (Not recommended in California).

Tomatoes, Eggplant, Peppers: Apply 4 to 6 weeks after transplanting. Plants should be well established and growing conditions favorable for good plant growth. Remove any weeds prior to application.

Precautions:

- . Do not use on beets, lettuce and other vegetables not listed on this label.
- . This product does not control established weeds.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING: Wear long-sleeved shirt and long-legged pants and chemical resistant gloves.

IMPORTANT: If the material comes in contact with skin, wash thoroughly with soap and water. Do not graze treated areas or feed plant parts to livestock. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting.

AFTER USE: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during use must be laundered separately from household articles.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply to a uniform weed-free soil surface within two to three days after removing any existing weeds. Disturbing the soil after application will reduce effectiveness.

This product does not control established weeds.

If there is no rain within 3 or 4 days after application, water for best results.

(Photo) Sprinkle the granules over the soil surface at the rate shown in this photo which equals 1 cup per 60 sq ft (6' x 10').

Flower Gardens: Apply after transplanting flowers and conditions are favorable for good plant growth. An additional application can be made anytime beds have been cultivated and are free of weeds. May be used around these flowers: Ageratum, alyssum, aster, babysbreath, bellflower, bleedingheart, bloodleaf, bugloss, candytufts, chrysanthemum, coleus, columbine, purple cornflower, coreopsis, coralbells, cosmos, cuphea, dahlia, delphinium, eveningprimrose, feverfew, forget-me-not, four-o'clock, foxglove, gaillardia, geranium, gladiolus, golddust, goldentuft, pink heath, iris, lantana, candle larkspur, lavendercotton, lily, lupine, golden marguerite, marigold, mourning bride, morningglory, rose moss, mother-of-thyme, nasturtium, orpine, peony, petunia, poker plant, rose, scarlet sage, snapdragon, Va. spiderwort, sedum, strawflower, sundrops, sunflower, sweet pea, African violet, wormwood, fernleaf yarrow, zinnia.

Shrubs, Trees and Groundcovers: Apply around newly transplanted or established plants after removing any existing weeds. May be used around these plants: Abella, Japanese andromeda, arborvitae, ash, azalea, barberry, birch, boxwood, camellia, chestnut, cinquefoil, cotoneaster, cottonwood, crabapple, cypress, deutzia, dogwood, elaeagnus, elm, euonymus, fir, forsythia, gum, hawthorn, holly, hydrangea, honeysuckle, Baltic ivy, Boston ivy, English ivy, Wilson ivy, juniper, lilac, locust, magnolia, maple, mockorange, mountain laurels, oak, pachistima, pachysandra, pine, pittosporum, podocarpus, poplar, privet, variegated privet, redbud, rhododendron, Russian olive, spirea, spruce, sycamore, tree peony, tulip tree, viburnum, walnut, weigela, willow, yew.

Strawberries (transplants or established beds): Apply in early spring prior to the first bloom. You may apply after harvest. Remove existing weeds prior to application.