

**XXXX**

Treats 2750 sq ft



**35-0-0**

# Bonus for dichondra

Weed control plus fertilizer for dichondra

### HOW TO USE

To control these actively growing weeds, apply any time and repeat in 8 weeks:

oxalis	filaree	soloiva
white clover	cudweed	poa annua
chickweeds	pearlwort	annual plantain

To prevent spotted spurge, apply in December, January or early February and repeat in 8 weeks.

To prevent crabgrass, apply in December, January, February or March and repeat in 8 weeks.

To ensure proper distribution use a Scott Spreader at setting 7

Slightly overlap spreader wheel tracks to provide complete coverage.

### PRECAUTIONS

Completely rinse particles off leaves immediately after application by adequate irrigation or hand hose spray.

Do not use on grass lawns.

On newly seeded areas wait 6 months before treating, on newly sodded or plugged areas wait one month. After application, wait 6 months before re-seeding.

Do not use on shaded areas receiving less than half a day of sunlight

This product does not control deep-rooted perennials such as bermuda grass, paspalums, dandelions and coarse fescue.

**Net Weight 12½ Lbs (5.7 Kg)**

ALL INFORMATION CONTAINED HEREIN IS THE PROPERTY OF SCOTT'S FERTILIZERS, INC. AND IS TO BE USED ONLY FOR THE PURPOSES SPECIFIED HEREIN. ANY REPRODUCTION OR TRANSMISSION OF THIS INFORMATION IN ANY FORM OR BY ANY MEANS WITHOUT THE WRITTEN PERMISSION OF SCOTT'S FERTILIZERS, INC. IS STRICTLY PROHIBITED. © 2000 SCOTT'S FERTILIZERS, INC.

**XXXX**



# 32-4-4 fertilizer and crabgrass preventer

FOR PROFESSIONAL USE ONLY  
for use on lawn and turfgrasses only

controls these weeds as they sprout  
while providing spring fertilization for turf

crabgrass	foxtail	lambquarters
barnyardgrass	goosegrass	pigweed

**SUGGESTED SPREADER SETTINGS**  
to provide proper distribution calibrate spreader before application

44 lbs (20.0 Kg) treats 10,000 sq ft at NORMAL RATE  
44 lbs (20.0 Kg) treats 8,000 sq ft at HEAVY RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			NORMAL RATE	HEAVY RATE
Scotts (Drop Type)	3 mph	overlap wheels	6 1/4	6 1/2
Scotts Rotary	3 mph	8 ft	H	J

\*For use only on large areas where there are no flower or garden seedbeds, flowers, shrubs or garden plants.

ACCEPTED  
March 12, 1974  
538-129  
Pesticide  
Label  
Subject  
U.S.

### DIRECTIONS FOR USE

To prevent crabgrass in the spring apply at NORMAL RATE before the third or fourth mowing of the year . . . before temperatures are regularly in the 80's.

To prevent barnyardgrass, foxtail, goosegrass, lambquarters and pigweed in addition to crabgrass apply at HEAVY RATE either in the fall (September-November) or in the spring.

Turf may be moist or dry.

### PRECAUTIONS

Do not seed, sod or sprig for four growing months following application.

On newly sodded or seeded areas, wait until grass has been mowed at least twice and is firmly rooted. On newly sprigged areas wait until bare areas are filled in.

After application, avoid any severe aerating or raking which might disturb the surface and break the chemical barrier.

Applications should not exceed twice yearly—allow at least a four month interval between applications.

Do not apply to flower or garden seedbeds, flowers, shrubs or garden plants.

ProTurf Division, O M Scott & Sons Co, Marysville, Ohio 43040

## Net Weight 44 Lbs (20.0 Kg)

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation and contact with skin, eyes or clothing. Wash after handling. Do not contaminate feed or foodstuffs or water used for irrigational or domestic purposes. Do not reuse container. Destroy when empty.

ENVIRONMENTAL CAUTIONS