

FRONT:


COVERS 500 SQ FT

(CONTENTS LIST)

STOP

Insects before They Start

Systemically control common sucking insects; aphids, leafhoppers, mites and scale on roses, shrubs and small trees (see back panel).



(Illustrations of aphids, leafhoppers, mites, or scale from list)

Systemic Action: Works from within the plant's own system. Stops sucking insects feeding on plants, even on new growth.

Soil Applied Granules. Not a spray or dust. Rain can't reduce product effectiveness.

(ILLUS.)  
( )  
( OF )  
( )  
( ROSE )

Protects new growth.

Protects hard-to-reach areas.

Work granules into soil.

Watering enables material to enter the root system.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes, and clothing. First aid: in case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. Avoid contamination of food and foodstuffs. This product may be a cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty container. Destroy when empty.

FRONT: (continued)

Required Clothing and Equipment for Application:

All applicators, including homeowners and all personnel involved with the mixing, loading, and transferring operations must wear the protective clothing and equipment enumerated below. The protective clothing and equipment to be worn is as follows:

- a. Impermeable gloves (for example, rubber or plastic covered reinforced gloves.)
- b. Boots or boot covers.
- c. Long-sleeved shirt and long pants.
- d. Wide-brimmed hat.
- e. Respirators must be worn by mixer/loaders.

ACTIVE INGREDIENT:

Dimethoate 0,0-dimethyl S-(methylcarbamoyl) methyl

phosphorodithioate . . . . . 5.00%

INERT INGREDIENTS: . . . . . 95.00%

Total 100.00%

Distributed by O.M. Scott & Sons Net Wt. 4 lbs. 10 oz. (c) 1982 O. M. Scott & Sons  
Marysville, OH 43041

LEFT SIDE

(Scotts (R))

Stop  
Insects Before They Start

SCOTTS NO QUIBBLE GUARANTEE

65168

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

Scotts (R)

Marysville, Ohio 43041

(Illustrations of aphids, leafhoppers, mites  
or scale from list; or plants from list)

RIGHT SIDE

(Scotts (R))

at  
Insects before they start

Scotts--dedicated to enhancing the natural beauty of our surroundings.

Use Scotts (R) STOP & GROW products for flower, shrub and tree care.

- . stop insects (liquid and spray)
- . stop disease (spray)
- . stop weeds . before they start
- . stop weeds . after they start
- . grow flowers
- . grow shrubs & trees
- . grow acid-loving plants
- . grow transplants

(Illustrations of aphids, leafhoppers, mites  
or scale; or plants from list)

Label

SCOTT'S (R)

Insect Control  
Insects Before They Start

ADVANTAGES OF SCOTT'S STOP INSECTS Before They Start:

Oil Soluble Form. Dry granules applied to the soil around plants with equipment and the avoidance of sprays and dust. Rain has no chance to wash off plant foliage or reduce effectiveness.

Systemic Action. Rain or watering enables roots to absorb the insecticide and circulate it through out the plant's own system. Stops sucking insects even on plant areas difficult to reach with sprays or dusts.

Long-lasting Results. Provides protection of the entire plant including new growth beginning 4-6 days after application and lasting 4-6 weeks.

Easy to Use. See below.

No Quibble Guarantee. See side panel

Directions for use.

Easy to use!

When: Starting in the spring or when insects first appear. Repeat every 4-6 weeks throughout the season or as needed.

Where: Systemically controls common sucking insects; Aphids on roses and elaeagnus; Leafhoppers on roses; Mites on roses, elaeagnus, poinsettia, pyracantha and viburnum; Scales on elaeagnus, poinsettia.

How: Loosen the surface soil beneath the spread of the branches (drip zone).

Apply product uniformly around base of plant using the tablespoons of product within the treatment width specified below.

Work product into the upper 2 inches of soil, so that granules are covered by one inch of soil.

<u>Plant Height</u> <u>(feet)</u>	<u>Tablespoons</u> <u>of Product</u>	<u>Treatment</u> <u>Widths</u> <u>(From base of plant)</u>
1	1	9 inches
2	2	13 inches
3	3	16 inches
4	5	20 inches
5	7	24 inches
6	9	28 inches

It is important to water in thoroughly immediately after application.  
Maintain good soil moisture (at least 1 inch per week) for 4-6 weeks  
after application.

PLEASE CONTINUE

Note: For plants that have been pulled, use the label, follow above instructions, then replace the mulch.

May be programmed separately with fertilizer products. Use the appropriate Scotts G.M. product.

May be programmed separately with products to control chewing insects (e.g., grasshoppers, caterpillars, beetles, etc.). Use Scotts STOP INSECT LIQUID or STOP INSECTS LIQUID or STOP INSECTS SPRAY.

May be programmed separately with products to control plant disease. Use Scotts STOP Disease Spray.

May be programmed separately with pre-emergence herbicide (Scotts STOP Weeds Before They Start). Apply insecticide prior to herbicide application since soil surface disturbance should be minimized for greater herbicide effectiveness.

Limitations:

Not a control for chewing insects (grasshoppers, caterpillars, beetles, etc.).

Do not use on spirea since plant injury may occur.

Do not use in the home or on potted plants or in vegetable gardens.

Store in a dry, ventilated area.

For Ground Application Only

EPA Reg. No. 538-135  
EPA Est. 34637-OH-1

O.H. Scott & Sons  
Marysville, Ohio 43041

Easy to Select Just What You Need for Flowers, Shrubs and Trees

(STOP & GROW line illustration)

TOP:

TO OPEN LEFT TAB

TO CLOSE:

FOLD LINER TIGHTLY

INSERT TAB HERE

(Scotts (A))

Stop  
Insects Before They Start