

**fertilizer plus weedgrass
preventer for bermudagrass**

NET WEIGHT 36 LBS (16.33 kg)

XXX



Scotts
ProTurf

30-4-4
fertilizer plus
weedgrass preventer
for bermudagrass

for use on common bermudagrass only

FOR PROFESSIONAL USE ONLY

NET WEIGHT 36 LBS (16.33 kg)

[illegible]

**fertilizer plus weedgrass
preventer for bermudagrass**

XXX

ACCEPTED
MAR 10 1989
U.S. Federal Insecticide,
Fungicide and Rodenticide Act,
and for the pesticide
C. A. King Inc.
registered under 538-143

30-4-4

**fertilizer plus
weedgrass preventer
for bermudagrass**

DIRECTIONS FOR USE

GENERAL CLASSIFICATION
It is a violation of Federal Law to use this product as
a means of communication with its labeling

Forest & common broadleaf tree species generation of these seeds	cedars evergreen spruce	Florida palm cedar	gum oak sycamore
--	-------------------------------	--------------------------	------------------------

SUGGESTED SPREADER SETTINGS

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	

For use only on common birdsong and only on projects
local that is as such as golf courses, golf links.

¹ For control of the *A. n. n.* it is necessary to apply to dry foliage in the middle of early spring, at grass is already growing, and immediately after application to remove ground granules from the foliage.

Wash seeds after the second moon. Before treating with seedbed cotton broadleafers, if things are wild, wash at least 4 weeks after longing. Do not seed or enter nursery with a treated area for at least one year.

Wash turf area level of leaves and more if necessary so that granules will reach the soil surface. Complete any raking or aerifying of the turf before treatment so the chemical barrier is against crabs and not the other weeds or not destroyed. Avoid contact with flowers and shrubs.

STORAGE AND DISPOSAL

It is a lightweight, portable, dry place out of reach of children and domestic animals. Close container tightly before storing. Do not store near water, feed or feedstuffs to be used for human or animal consumption.

Do not reuse container. Dispose of it as waste of an unused amount for another container. Do not use a safe.

Protest Division 08 3rd & 3rd, Maryland Ave 4 1960

**fertilizer plus weedgrass
preventer for bermudagrass**

NET WEIGHT 36 LBS (16.33 kg)

XXX



ProTurf

**30-4-4
fertilizer plus
weedgrass preventer
for bermudagrass**

for use on common bermudagrass only

FOR PROFESSIONAL USE ONLY

NET WEIGHT 36 LBS (16.33 kg)

[illegible]

XXXX

ACCEPTED

MAR 10 1989

Under Federal Insecticide,
Fungicide and Rodenticide Act,
Registration No. 538-143
Registered under

30-4.4

**fertilizer plus
weedgrass preventer
for bermudagrass**

DEFINITIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Further research demonstrates, prevents germination of these seeds:

hairy grass	covered seeds	hard seed	strongly spotted
crabgrass	protrude	trial	spruce

SUGGESTED COMMAND SETTINGS

[illegible]

For use only on cotton in Mississippi and only on professional turf areas such as golf courses, and farms.

For control of the annual weeds listed above apply to dry foliage on the soil or early spring. If grass is actively growing, water immediately after application to remove product granules from the foliage.

Wait until after the second mowing before treating newly seeded common bermudagrass; if sprays are used, wait at least 4 weeks after sprigging. Do not use or make overseed into a treated area for at least one year.

With turf area free of leaves and once if necessary so that ground will reach the soil surface. Complete any raising or leveling of the turf before treatment in the chemical burn-off against crabgrass and the other weeds is not destroyed avoid contact with flowers and shrubs.

STORAGE AND DISPOSAL

Store in original container in a dry place out of reach of children and domestic animals. Close container tightly before storing. Do not store near water, feed or feedstuffs to be used for human or animal consumption.

Do not reuse container. Dispose of in an incinerator or landfill approved for pesticide containers or burn in a safe stove.

Proctor Division, O M Scott & Son - Maryland, Ohio 43540

**fertilizer plus weedgrass
prevents for bermudagrass**

XXX

38