

CON PROFESSIONAL USE ONE TO A CONTROL OF THE PROFESSIONAL USE ONE OF THE PROFESSIONAL USE ONE OF THE PROFESSIONAL USE OF THE P



DIRECTIONS FOR USE

for use on summer turf diseases

Fertilizes turfgrasses. Controls fungus activity causing brownpatch, leafspot, dollarspot, red thread and copperspot. Also controls moss.

SUGGESTED SPREADER SETTINGS

To provide proper distribution calibrate spreader before application.

46½ LBS (21.1 Kg) TREATS 22,000 SQ FT (½ ACRE/2044 SQ M) AT NORMAL RATE 46½ LBS (21.1 Kg) TREATS 11,000 SQ FT (¼ ACRE/1022 SQ M) AT DOUBLE RATE 46½ LBS (21.1 Kg) TREATS 44,000 SQ FT (1 ACRE/4088 SQ M) AT HALF RATE

| SPREADER | GROUND OR PTO SPEED | WIDTH OF COVERAGE | SPREA NORMAL RATE | DER SETT OUUBLE RATE | TING HALF RAYE |
|---------------------------------|------------------------|----------------------|-------------------------|----------------------------|----------------------|
| Scotts (Drop Type) | | overlap wheels | 4 1/2 D | 6 G | 31/2 B |
| Scotts Rotary Lely-WTR & | 3 mph | | _ | | |
| WFR Models Lely-HR | 4.5 mph 450 rpm | 16 feet 16 feet | | 31/2 IV 31/2 IV | |
| (PTO Model) Gandy 8 ft A Series | @4.5 mph 4.5 mph | overlap wheels | 131/2 | 171/2 | 10% |
| Model 42 model | 2.5 mph | overlap wheels | 24 | 30 | 19 |

Apply to moist or dry grass.

Putting greens—may be used every two weeks at NORMAL RATE during growing season for prevention or control. Interim protection can be achieved with ProTurf Broad Spectrum Fungicide. Syringe surface lightly after application to avoid pickup on shoes, golf balls or maintenance equipment.

All other areas—use before fungus damage appears or when symptoms are first noticed. May be repeated every month at NORMAL RATE. If disease is severe or more feeding is desired apply at DOUBLE RATE.

To control moss—apply anytime at NORMAL RATE to moist foliage. Follow up at two week intervals as needed.

ProTurf Division, OM Scott & Sons, Marysville, Ohio 43040