


ST AUGUSTINE WEED CONTROL
PLUS FERTILIZER (S-1207)
XXXXX

FRONT:

03/28/84 RF



XXXXX

(Stock number will be inverted.)

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

- . controls weeds in St Augustine, zoysia, centipede and carpetgrass
- . provides controlled release nitrogen feeding

Net Weight 55 lbs (24.95 kg)

LEFT COLUMN

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

Environmental Hazards: On compact, slowly draining soils, (particularly the Dallas, Fort Worth area), avoid application on rootzone of trees and shrubs.

ACTIVE INGREDIENT:

Atrazine, 2-Chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine		1.32%
INERT INGREDIENTS:		99.68%
EPA Reg No 538-157	EPA Est 538-OH-1	Total 100.00%

FRONT CONTINUED:

RIGHT COLUMN

St Augustine Weed Control Plus Fertilizer 26-3-3 643-XXXX

Guaranteed Analysis

Total nitrogen (N)	26%
0.7% ammoniacal nitrogen	
8.0% methylenediurea and dimethylenetriurea nitrogen	
16.0% urea, other methylene ureas nitrogen	
1.3% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	3%
Derived from monoammonium phosphate, urea, methylene ureas and muriate of potash.	
Chlorine, not more than	2.4%
Product of USA	

ProTurf Division, The O M Scott & Sons Company Marysville, Ohio 43041
(c)1985, The O M Scott & Sons Company. All rights reserved.

For Professional Use Only

XXXXX

GUSSETS:

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine Weed
Control Plus Fertilizer

LEFT GUSSET

(
(
Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine Weed
Control Plus Fertilizer

XXXXX

BACK:

XXXXX

(Stock number will be inverted.)

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine Weed
Control Plus Fertilizer

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls weeds listed below plus fertilizes the turf.

Apply to control the following weeds any time they are growing. If complete control is not obtained a second treatment may be required 2 months after the first.

annual sowthistle
black medic
blue toadflax
buckhorn plantain
burclover
Carolina geranium
(cranesbill)
common chickweed
common dandelion
common purslane
cutleaf
 eveningprimrose
dichondra
dollarweed
Florida betony

Florida pusley
fragrant cudweed
 (rabbit-tobacco)
henbit
lippia (matchweed)
oxalis (common
 yellow woodsorrel)
red sorrel
rescuegrass
shepherdspurse
stinging nettle
Venus lookingglass
Virginia pepperweed
wild mustard

Apply in the fall to control winter weeds such as henbit, chickweed, rabbit-tobacco, and eveningprimrose.

Apply in January, February or early March to prevent germination of the following annual weeds. Product does not control established plants of these weeds.

crabgrass
bull paspalum

sedges (from seed)
spotted spurge

Make all applications to dry foliage only. If maximum daytime temperature is 80° or above. Water sufficiently to rinse particles off the grass blades.

Application can be made to newly sprigged St Augustine and zoysia turf. Prior to this treatment, however, water should be applied to settle the soil around the sprigs.

BACK CONTINUED:

Use limitations

Do not apply to lawns of bahia, bluegrass, bentgrass, dichondra, ryegrass, tall fescue, bermudagrass or mixed lawns containing any of these grasses.

Do not use on centipede or carpetgrass turf during the period of winter dormancy (approximately October 15 to March 15), except in Florida.

Do not apply to newly seeded zoysia, centipede or carpetgrass until after the turf has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product.

Do not apply on bedding plants, flowers, garden plants or ornamental trees and shrubs.

Where winter overseeding is planned, do not treat the turf during the preceding 6 months. Application the following spring hastens conversion to the summer turf while controlling weeds and feeding the grass.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

BACK CONTINUED:

Suggested Spreader Settings
to provide proper distribution calibrate spreader before application

55 lbs (24,95 kg) treats 11,000 ft² (1/4 acre/1022 m²) at RECOMMENDED RATE

<u>SPREADER</u>	<u>GROUND PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTING RECOMMENDED RATE</u>
Scotts (Drop Type)	3 mph		
Scotts Rotaries	3 mph		
Lely - WTR Model	4.5 mph		
Lely - HR	450 mph		
(PTO Model)	4.5 mph		

XXXXX

26-3-3
St Augustine
Weed Control
Plus Fertilizer