

538-157

PA125

142

84798

26-3-3
St Augustine
Weed Control
Plus Fertilizer

ACCEPTED
AUG 19 1987
U.S. Department of Agriculture
Fungicide, Herbicide, Insecticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 538-157



26-3-3 St Augustine Weed Control Plus Fertilizer

6/5/86



26-3-3
St Augustine
Weed Control
Plus Fertilizer

26-3-3
St Augustine
Weed Control
Plus Fertilizer



- controls weeds in St Augustine, zoysia, centipede and carpetgrass
- provides controlled release nitrogen feeding

Net Weight 50¹³/₁₆ lbs (23.05 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from food or food products.

Environmental Hazards: On compact, slowly draining soils (particularly the Dallas Fort Worth area) avoid application on rootzone of trees and shrubs. Airborne residues readily and accepted leaf rates have been found to result in contamination of water supplies by way of groundwater. Therefore users are advised to avoid use of atrazine in well drained soils particularly in areas having high groundwater tables. All crops - This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENT
Atrazine (2-Chloro-4-ethylamino-6-isopropylamino-s-triazine) 0.99%
INERT INGREDIENTS 99.01%
Total 100.00%

EPA Reg No 538-157 EPA Est 538-04-1
St Augustine Weed Control Plus Fertilizer 26-3-3
Oxamethozon Analyses

643-8548

Total nitrogen (N)	26%
0.6% ammoniacal nitrogen	
24.1% urea methylene urea nitrogen*	
1.3% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	7%
Soluble potash (K ₂ O)	
Derived from monoammonium phosphate urea methylene urea and the rate of potash	
*Conc. a 8.1% slowly available methylenediamine and dimethylenediamine nitrogen	
Chlorine not more than	2.5%
Product of USA	
U.S. Pat No 4 378 238	

ProLurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041
© 1987 The O M Scott & Sons Company. All rights reserved.

84798

84798

Recommended for Use by Professional Turfgrass Managers

2062



26-3-3 St Augustine Weed Control Plus Fertilizer

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls weeds listed below plus fertilizes the turf.

Apply to control the following weeds any time they are growing. If complete control is not obtained a second treatment may be required 2 months after the first.

- | | |
|-------------------|---------------------|
| annual sowthistle | Florida pusley |
| black medic | fragrant cutweed |
| blue iceplant | (rabbit-tobacco) |
| buckhorn plantain | herb |
| burclover | lupine (marchweed) |
| Carolina geranium | oxalis (common) |
| (cransbill) | yellow woodsorrel) |
| common chickweed | red sorrel |
| common dandelion | rescuegrass |
| common purslane | shepherdspurse |
| cutleaf | stinging nettle |
| eveningprimrose | Venus lookingglass |
| dichondra | Virginia peppercorn |
| dollarweed | wild mustard |
| Florida betony | |

Apply in the fall to control winter weeds such as hen bit, chickweed, rabbit-tobacco, and eveningprimrose.

Apply in January, February or early March to prevent germination of the following annual weeds. Product does not control established plants of these weeds.

- | | |
|----------------------|--------------------|
| crabgrass | sedges (from seed) |
| (Digitaria serotina) | spotted spurge |
| bull paspalum | |

Make all applications to dry foliage only. If maximum daytime temperature is 80°F or above, water sufficiently to rinse particles off the grass blades.

Application can be made to newly sprigged St Augustine and zoysia turf. Prior to this treatment, however, water should be applied to settle the soil around the sprigs.

Use limitations

Do not apply to lawns of bahia, bluegrass, bentgrass, dichondra, ryegrass, tall fescue, bermudagrass or mixed lawns containing any of these grasses.

Do not use on centipede or carpetgrass turf during the period of winter dormancy (approximately October 15 to March 15), except in Florida.

Do not apply to newly seeded zoysia, centipede or carpetgrass until after the turf has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product.

Do not apply on bedding plants, flowers, garden plants or ornamental trees and shrubs.

Where winter overseeding is planned, do not treat the turf during the preceding 6 months. Application the following spring hastens conversion to the summer turf while controlling weeds and feeding the grass.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings			
to provide proper distribution calibrate spreader before application.			
30" x 1/2" lbs treats 11,000 ft ² (1/4 acre) at the RECOMMENDED RATE			
SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERED overlap wheels	RECOMMENDED RATE
Scotts Drop	3 mph		7
Scotts Backpack			
R-7, R-7T	3 mph	8 feet	1 1/4
R-6 (cone B)	3 mph	8 feet	M
Lady Models (See below)	4.5 mph	28 feet	6 M

Pattern shown to the left. Pattern to the right is for the right.
Lady models R-7, R-7T, R-6 and R-6T are 1200 lbs. at 400 gals.
Make settings 20-25 lbs. per acre 1000 or at the 1000 lb. recommended rate.

26-3-3
St Augustine
Weed Control
Plus Fertilizer

84798