

538-157 PM25 10612
PROTURF ST AUGUSTINE WEED CONTROL
PLUS FERTILIZER (MURP) XXXXX
04/13/88 CR/VS

(Scotts(R))
ProTurf(R)
BRAND


26-3-3
St Augustine
Weed
Control Plus
Fertilizer

- o controls weeds in St Augustine, zoysia, centipede and carpetgrass
 - o provides controlled release nitrogen feeding
-

Net Weight 55 lbs (24.95 kg)

LEFT COLUMN

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS



Hazards to Humans and Domestic Animals: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

Environmental Hazards: On compact, slowly draining soils, (particularly the Dallas, Fort Worth area), avoid application on rootzone of trees and shrubs. Atrazine leaches readily and accepted label rates have been found to result in contamination of water supplies by way of groundwater. Therefore users are advised to avoid use of atrazine in well drained soils particularly in areas having high groundwater tables. All crops--This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENT:	
Atrazine (2-Chloro-4-ethylamino-6-isopropylamino-S-triazine) . . .	1.32%
INERT INGREDIENTS:	<u>98.68%</u>
	Total 100.00%

RIGHT COLUMN

EPA Reg. No. 538-157	EPA Est 538-OH-1	
St Augustine Weed Control Plus Fertilizer	26-3-3	643-XXXX

Guaranteed Analysis

Total nitrogen (N)	26%
0.7% ammoniacal nitrogen	
24.0% urea, methylene ureas nitrogen*	
1.3% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	3%
Derived from: monoammonium phosphate, urea, methylene ureas and muriate of potash.	

* Contains X.X% slowly available methylenediurea and dimethylenetriurea nitrogen.

Chlorine, not more than. 2.4%

Product of USA

U.S. Pat No XXXXXXXXX

ProTurf Division, The O M Scott & Sons Company
 Marysville, Ohio 43041
 (c) 1988, The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

XXXXX

26-3-3
 St Augustine
 Weed Control
 Plus Fertilizer

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

XXXXX

XXXXX

400 12
26-3-3
St Augustine
Weed Control
Plus Fertilizer

(NOTE: The description above will be inverted.)

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls weeds listed below plus fertilizes the turf.

Apply to control the following weeds any time they are growing. If complete control is not obtained a second treatment may be required 2 months after the first.

- | | |
|-------------------|---------------------|
| annual sowthistle | Florida pusley |
| black medic | fragrant cudweed |
| blue toadflax | (rabbit-tobacco) |
| buckhorn plantain | henbit |
| burclove | lippia (matchweed) |
| Carolina geranium | oxalis (common) |
| (craneshill) | yellow woodsorrel) |
| common chickweed | red sorrel |
| common dandelion | rescuegrass |
| common purslane | shepherdspurse |
| cutleaf | stinging nettle |
| eveningprimrose | Venus lookingglass |
| dichondra | Virginia pepperweed |
| dollarweed | wild mustard |
| Florida betony | |

Apply in the fall to control winter weeds such as henbit, chickweed, rabbit-tobacco, and eveningprimrose.

Apply in January, February or early March to prevent germination of the following annual weeds. Product does not control established plants of these weeds.

- | | |
|----------------------|--------------------|
| crabgrass | sedges (from seed) |
| (Digitaria serotina) | spotted spurge |
| bull paspalum | |

Make all applications to dry foliage only. If maximum daytime temperature is 80°F or above, water sufficiently to rinse particles off the grass blades.

Application can be made to newly sprigged St Augustine and zoysia turf. Prior to this treatment, however, water should be applied to settle the soil around the sprigs.

RIGHT COLUMN

Use limitations

Do not apply to lawns of bahia, bluegrass, bentgrass, dichondra, ryegrass, tall fescue, bermudagrass or mixed lawns containing any of these grasses.

Do not use on centipede or carpetgrass turf during the period of winter dormancy (approximately October 15 to March 15), except in Florida.

Do not apply to newly seeded zoysia, centipede or carpetgrass until after the turf has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product.

Do not apply on bedding plants, flowers, garden plants or ornamental trees and shrubs.

Where winter overseeding is planned, do not treat the turf during the preceding 6 months. Application the following spring hastens conversion to the summer turf while controlling weeds and feeding the grass.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings
to provide proper distribution calibrate spreader before application.

55 lbs treats 11,000 ft² (1/4 acre) at the RECOMMENDED RATE

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE</u>
Scotts Drop	3 mph	X	X
Scotts Rotaries			
R-7, R-7X*	3 mph	X feet	X
R-8 (cone 9)	3 mph	X feet	X
Lely Models (See below)	4.5 mph	XX feet	X

*Pattern skews to the left. Pattern is 4 1/2 ft to the left, 3 1/2 ft to the right.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.
Metric coverage: 24.95 kg treats 1022 m² at the RECOMMENDED RATE.

LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O. M. Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O. M. Scott & Sons Company.

ProTurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041

(Scotts(R))
ProTurf(R)
BRAND


26-3-3
St Augustine
Weed
Control Plus
Fertilizer

- o controls weeds in St Augustine, zoysia, centipede and carpetgrass
 - o provides controlled release nitrogen feeding
-

Net Weight 63 1/2 lbs (20.80 kg)

LEFT COLUMN

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS



Hazards to Humans and Domestic Animals: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

Environmental Hazards: On compact, slowly draining soils, (particularly the Dallas, Fort Worth area), avoid application on rootzone of trees and shrubs. Atrazine leaches readily and accepted label rates have been found to result in contamination of water supplies by way of groundwater. Therefore users are advised to avoid use of atrazine in well drained soils particularly in areas having high groundwater tables. All crops--This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

GUSSETS:

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

XXXXX

XXXXX

26-3-3
St Augustine
Weed Control
Plus Fertilizer

(NOTE: The description above will be inverted.)

(Scotts(R))
ProTurf(R)
BRAND

26-3-3
St Augustine
Weed Control
Plus Fertilizer

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls weeds listed below plus fertilizes the turf.

Apply to control the following weeds any time they are growing. If complete control is not obtained a second treatment may be required 2 months after the first.

- | | |
|-------------------|---------------------|
| annual sowthistle | Florida pusley |
| black medic | fragrant cudweed |
| blue toadflax | (rabbit-tobacco) |
| buckhorn plantain | henbit |
| burclover | lippia (matchweed) |
| Carolina geranium | oxalis (common |
| (cranesbill) | yellow woodsorrel) |
| common chickweed | red sorrel |
| common dandelion | rescuegrass |
| common purslane | shepherdspurse |
| cutleaf | stinging nettle |
| eveningprimrose | Venus lookingglass |
| dichondra | Virginia pepperweed |
| dollarweed | wild mustard |
| Florida betony | |

Apply in the fall to control winter weeds such as henbit, chickweed, rabbit-tobacco, and eveningprimrose.

Apply in January, February or early March to prevent germination of the following annual weeds. Product does not control established plants of these weeds.

- | | |
|----------------------|--------------------|
| crabgrass | sedges (from seed) |
| (Digitaria serotina) | spotted spurge |
| bull paspalum | |

Make all applications to dry foliage only. If maximum daytime temperature is 80°F or above, water sufficiently to rinse particles off the grass blades.

Application can be made to newly sprigged St Augustine and zoysia turf. Prior to this treatment, however, water should be applied to settle the soil around the sprigs.

RIGHT COLUMN

Use limitations

Do not apply to lawns of bahia, bluegrass, bentgrass, dichondra, ryegrass, tall fescue, bermudagrass or mixed lawns containing any of these grasses.

Do not use on centipede or carpetgrass turf during the period of winter dormancy (approximately October 15 to March 15), except in Florida.

Do not apply to newly seeded zoysia, centipede or carpetgrass until after the turf has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product.

Do not apply on bedding plants, flowers, garden plants or ornamental trees and shrubs.

Where winter overseeding is planned, do not treat the turf during the preceding 6 months. Application the following spring hastens conversion to the summer turf while controlling weeds and feeding the grass.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings
to provide proper distribution calibrate spreader before application.

63 1/2 lbs treats 11,000 ft² (1/4 acre) at the RECOMMENDED RATE

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE</u>
Scotts Drop	3 mph	overlap wheels	X
Scotts Rotaries			
R-7, R-7X*	3 mph	X feet	X
R-8 (cone 9)	3 mph	X feet	X
Lely Models	4.5 mph	XX feet	X
(See below)			

*Pattern skews to the left. Pattern is 4 1/2 ft to the left, 3 1/2 ft to the right.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.
Metric coverage: 20.80 kg treats 1022 m² at the RECOMMENDED RATE.

LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O. M. Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O. M. Scott & Sons Company.

ProTurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041