


3/8" →



29-3-4 St. Augustine Weed Control Plus Fertilizer

8664-6



# 29-3-4 St. Augustine Weed Control Plus Fertilizer

- Controls weeds in St. Augustine, zoysia, centipede and carpetgrass
- Provides controlled release nitrogen feeding

**ACCEPTED**  
 OCT 30 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 538-157

### KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

Hazardous to Marine and Benthic Animals  
Harmful if swallowed. Avoid inhalation or contact with eyes. Keep away from food or food products.

**Personal Protective Equipment**  
Applicator and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

#### Use Safety Recommendations

Users should:

- Wash hands before eating, drinking, smoking, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside; then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards:** On compact, slowly draining soils, (particularly the Dallas, Fort Worth area), avoid application on receptors of trees and shrubs. Airborne leaches readily and significant leach rates have been found to result in contamination of water supplies by use of granulars. Therefore, users are advised to avoid use of granules in well drained soils particularly in areas having high groundwater tables. All granules—this pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

#### ACTIVE INGREDIENT:

Atrazine	1.055%
RELATED COMPOUNDS:	0.055%
OTHER INGREDIENTS:	98.89%
<b>Total</b>	<b>100.00%</b>

TCAS #1912-24-9

EPA Reg. No. 538-157

EPA Est. 538-OH-151  
39578-TX-1E1

Subscript used is first letter of run code on end of container.

St. Augustine Weed Control Plus Fertilizer 29-3-4 F 643

#### Guaranteed Analysis

Total nitrogen (N)	29%
6.2% ammoniacal nitrogen	
12.3% urea nitrogen	
9.1% other water soluble nitrogen	
0.9% water insoluble nitrogen	
Available phosphate (P <sub>2</sub> O <sub>5</sub> )	3%
Soluble potash (K <sub>2</sub> O)	4%
Sulfur (S)	8.5%
0.5% combined sulfur (S)	

Derived from urea, methylene urea, ammonium sulfate, ammoniacal phosphate and potassium sulfate.  
\*Contains 6.7% slowly available methylenediamine and dimethylenediamine nitrogen.

Product of U.S.A.  
U.S. Pat. No. 4,379,238 and 5,102,460

Professional Business Group  
The Scotts Company  
14111 Swanton Rd.  
Marysville, OH 43041

©1998, The Scotts Company. World rights reserved.

## Net Weight 45.52 lb (20.64 kg)

Recommended For Use By Professional Turfgrass Managers



VID-80  
15175R0  
47483  
10/98



29-3-4 St. Augustine Weed Control Plus Fertilizer

8664-6

3/8"



29-3-4 St. Augustine Weed Control Plus Fertilizer 8664-6

29-3-4 St. Augustine Weed Control Plus Fertilizers



CUSTOMER INK JET AREA

3/8"

1 1/2 2/2

# 29-3-4 St. Augustine Weed Control Plus Fertilizer

## Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, waterproof gloves, shoes plus socks.

Controls weeds listed below plus fertilizes the turf.

Apply to control the following weeds any time they are growing. If complete control is not obtained, a second treatment may be required 2 months after the first.

- annual sowthistle
- black medic
- blue joadite
- buckhorn plantain
- burdock
- Carolina geranium (cranesbill)
- common chickweed
- common dandelion
- common purslane
- cutleaf eveningprimrose
- dichondra
- dollarweed
- Florida betony
- Florida pusley
- fragrant eyebeard (fish-tail tobacco)
- herb
- hippis (mitchweed)
- oasis (common yellow wood-saw)
- red sorrel
- rescuegrass
- shepherdspurse
- string nettle
- Verus looksgloss
- Virginia pepperweed
- wild mustard

Apply in the fall to control winter weeds such as herb, chickweed, rabbit-tobacco, and eveningprimrose.

Apply in January, February or early March to prevent germination of the following annual weeds. Product does not control established plants of these weeds.

- craggrass (Digitaria serotina)
- bulr-paspalum
- sedges (from seed)
- spotted spurge

Make all applications to dry foliage only, if maximum daytime temperature is 80°F or above, water sufficiently to rinse particles off the grass blades.

Application can be made to newly sprigged St. Augustine and zoysia turf. Prior to this treatment, however, water should be applied to settle the soil around the sprigs.

### Use limitations

Do not apply to lawns of bahia, bluegrass, bentgrass, dichondra, ryegrass, tall fescue, bermudagrass or mixed lawns containing any of these grasses.

Do not use on centipede or carpetgrass turf during the period of winter dormancy (approximately October 15 to March 15), except in Florida.

Do not apply to newly seeded zoysia, centipede or carpetgrass until after the turf has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product.

Do not apply on bedding plants, flowers, garden plants or ornamental trees and shrubs.

Where winter overseeding is planned, do not treat the turf during the preceding 6 months. Application the following spring hastens conversion to the summer turf while controlling weeds and feeding the grass.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its use based on the label. The user should always follow the label directions and exercise his judgment and caution when using the product under the growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or buyer, and the limit of liability of the Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and user are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.



Professional Business Group  
The Scotts Company  
14111 Scottslawn Rd.  
Marysville, OH 43041  
www.scottscompany.com

### Suggested Spreader Settings

To provide proper distribution, calibrate your spreader before application.

45.52 lb treats 11,000 sq. ft. (1/4 acre) at the Recommended Rate

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE
Scotts Drop Type	3 mph	overlap wheels	6 1/2
Scotts Rotaries R-7, R-7X*	3 mph	11 feet	I
R-8, R-8A (Cone 9)	3 mph	11 feet	L
AzcuPro, SR-2000 (Cone 9)	3 mph	11 feet	N
Gandy (30")	3 mph	overlap wheels	29
Vicon (Q3 Series - Standard Spout)	4.5 mph	24 feet	26
Lady Models WTR, WFR, HR, 1250. (PTO at 450 rpm)	4.5 mph	30 feet	6 1/4 III

\* Slide 1/4 closed.

SEAL AREA (NO PRINT)

EDGE OF

1 1/2" 1" 1/2"