



Fungicide VI

8848

Fungicide VI

• prevents and controls turfgrass diseases including benzimidazole tolerant dollar spot

Net Weight 23 lbs (10.44 kg)

BEST DOCUMENT AVAILABLE

For Professional Use Only

8848 Fungicide VI

Fungicide VI



8848

35

Fungicide VI 8848



Fungicide VI

Directions for use

For a complete list of turfgrass diseases to which this product is effective, see the product label.

Control the following turfgrass diseases: Dollar spot, Brown patch, Red leaf spot, Fusarium patch (sheep scurf), and Leaf spot (including benzimidazole tolerant dollar spot).

DOLLAR SPOT, BROWN PATCH and RED LEAF SPOT: To prevent disease apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

For active disease apply to moist foliage at the DOUBLE RATE and lightly water or irrigate shortly to remove some of the product particles from the foliage. Repeat the application every 7 weeks. Once the disease is under control use the RECOMMENDED RATE applications at 2 week intervals for the remainder of the disease period.

FUSARIUM PATCH (Sheep Scurf and Cal) only: To prevent disease apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 to 3 week intervals throughout the expected disease period.

For active Fusarium patch apply to moist foliage at the DOUBLE RATE and repeat at 2 to 3 week intervals. When the disease activity is under control, apply at the RECOMMENDED RATE at 2 to 3 week intervals.

LEAF SPOT (including Sheepshead and Cal) only: To prevent disease apply to moist foliage at the DOUBLE RATE when environmental conditions

favor disease activity. Repeat if needed after 3-4 weeks. For active leaf spot apply to moist foliage at the DOUBLE RATE when disease activity is first noted. Repeat if needed at the DOUBLE RATE at intervals not closer than 2 weeks.

Reminder

Make all applications to moist foliage. At DOUBLE RATE applications on putting green turf, except for Fusarium patch and leaf spot, should be lightly watered or irrigated shortly after application. Heavy watering may significantly reduce disease control.

Do not mow for 24 hours after application. This product is to be applied with drop spreaders only.

Please pay special attention to the handling instructions as noted on the warning statement to ensure safety of the product.

Storage and disposal

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Pesticides that cannot be used should be disposed of in a unit approved for pesticides or buried in a safe place away from water supplies.

Do not reuse empty container. Dispose of in an incinerator or landfill approved for pesticides or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as incinerated open burning.

ProLurf Division, D M Scott & Sons, Marysville, Ohio 43040

Suggested Spreader Settings	
to prevent proper disease control, check spreader before application.	
23 lbs (10.44 kg) spreader	11.000 ft (3.353 m) at RECOMMENDED RATE
23 lbs (10.44 kg) spreader	11.000 ft (3.353 m) at DOUBLE RATE
23 lbs (10.44 kg) spreader	11.000 ft (3.353 m) at DOUBLE RATE

3

35



Fungicide VI 8848



Fungicide VI



Fungicide VI

8848

Fungicide VI

• prevents and controls turfgrass diseases including benzimidazole tolerant dollar spot

Net Weight 23 lbs (10.44 kg)

BEST DOCUMENT AVAILABLE

For Professional Use Only

8848 Fungicide VI

Fungicide VI



8848

35

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Consult the following turfgrass diseases:

Dollar spot, Red leaf spot, Brown patch, Fusarium patch (Wash state), and other diseases.

For active dollar spot apply to most foliage at the DOUBLE RATE and lightly water immediately to remove some of the product particles from the foliage. Repeat the application every 2 weeks if the disease is under control use RECOMMENDED RATE applications at 2 week intervals for the remainder of the disease period.

For active Fusarium patch apply to most foliage at the DOUBLE RATE and repeat at 2 to 3 week intervals when the disease activity is under control apply at the RECOMMENDED RATE at 2 to 3 week intervals.

LEAF SPOT (Microdochium spp.) To prevent dollar spot apply to most foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat the application at 2 to 3 week intervals throughout the expected disease period.

For active leaf spot apply to most foliage at the DOUBLE RATE when disease activity is first noted. Repeats needed at the DOUBLE RATE at intervals that closely meet 2 weeks.

Reminder

Make all applications to most foliage at the DOUBLE RATE applications on cutting grass turf except for Fusarium patch and leaf spot should be lightly watered immediately after application. Heavy watering may significantly reduce disease control. Delay watering for 24 hours after application. This product is to be applied with drop spreaders only. Please pay special attention to the following information as noted in the warning statement to ensure safe use of the product.

Storage and disposal

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Pesticides that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse empty containers. Dispose of as an empty container or in a safe place. Consult Federal, State or local disposal authorities for approved container procedures such as incineration or open burning.

© 1987 Division of M. Scott & Sons, Moline, Ohio 43060

Suggested Spreader Settings	
1/2" to 3/4" spreader	1000
3/4" to 1" spreader	1200
1" to 1 1/2" spreader	1400
1 1/2" to 2" spreader	1600
2" to 3" spreader	1800
3" to 4" spreader	2000
4" to 6" spreader	2200
6" to 8" spreader	2400
8" to 10" spreader	2600
10" to 12" spreader	2800
12" to 14" spreader	3000
14" to 16" spreader	3200
16" to 18" spreader	3400
18" to 20" spreader	3600
20" to 22" spreader	3800
22" to 24" spreader	4000
24" to 26" spreader	4200
26" to 28" spreader	4400
28" to 30" spreader	4600
30" to 32" spreader	4800
32" to 34" spreader	5000
34" to 36" spreader	5200
36" to 38" spreader	5400
38" to 40" spreader	5600
40" to 42" spreader	5800
42" to 44" spreader	6000
44" to 46" spreader	6200
46" to 48" spreader	6400
48" to 50" spreader	6600
50" to 52" spreader	6800
52" to 54" spreader	7000
54" to 56" spreader	7200
56" to 58" spreader	7400
58" to 60" spreader	7600
60" to 62" spreader	7800
62" to 64" spreader	8000
64" to 66" spreader	8200
66" to 68" spreader	8400
68" to 70" spreader	8600
70" to 72" spreader	8800
72" to 74" spreader	9000
74" to 76" spreader	9200
76" to 78" spreader	9400
78" to 80" spreader	9600
80" to 82" spreader	9800
82" to 84" spreader	10000
84" to 86" spreader	10200
86" to 88" spreader	10400
88" to 90" spreader	10600
90" to 92" spreader	10800
92" to 94" spreader	11000
94" to 96" spreader	11200
96" to 98" spreader	11400
98" to 100" spreader	11600

35A