




Fungicide VI

8848



Fungicide VI

prevents and controls turfgrass diseases including benzimidazole tolerant dollar spot

Net Weight 23 lbs (10.44 kg)

For Professional Use Only

8848

Fungicide VI

Fungicide VI



8848

Fungicide VI 8848



Fungicide VI

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls the following turfgrass diseases:
brown patch, dollar spot, *Heterobasidium* spp. (including red leaf spot), *benzimidazole*, *Fusarium* patch (leaves), *Stemmatium* (leaves), *Doth* and *Colt* only.

DOLLAR SPOT, BROWN PATCH and RED LEAF SPOT: To prevent disease apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the spot control at 2 week intervals throughout the reported disease period.

For active disease apply to moist foliage at the DOUBLE RATE and irrigate water immediately to remove some of the product particles from the foliage. Repeat the application every 2 weeks. Once the disease is under control use RECOMMENDED RATE applications at 2 week intervals for the remainder of the disease period.

FUSARIUM PATCH (leaves), Doth and Colt only: To prevent disease apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 to 3 week intervals throughout the reported disease period.

For active *Fusarium* patch apply to moist foliage at the DOUBLE RATE and repeat at 2 to 3 week intervals. When the disease activity is under control apply at the RECOMMENDED RATE at 2 to 3 week intervals.

LEAF SPOT (*Heterobasidium* spp.): To prevent disease apply to moist foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat if needed after 3-4 weeks.

For active leaf spot apply to moist foliage at the DOUBLE RATE when disease activity is first noted. Repeat as needed at the DOUBLE RATE at intervals not longer than 2 weeks.

Reminder

Make all applications to moist foliage.

At DOUBLE RATE applications to prevent disease, but except for Fungicide VI, applications should be light, evened or immediately after application where watering may apply only once or twice a week.

Delay mowing for 24 hours after application. This product is to be applied with drop spreaders only.

Do not graze treated areas. Do not feed. Observe the label's instructions and keep off treated areas until mowers have been washed off the foliage and the area is completely dry.

Please pay special attention to the handling instructions as noted in the warning statement to ensure safe use of the product.

Storage and disposal

STORAGE: Store in a clean, dry place. Remove opened bag by folding top down and securing.

RESIDUE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Residue resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a secondary landfill or by incineration as it allowed by State and local authorities. Do not burn. If burned, stay out of smoke.

LIMITATION OF LIABILITY: It must be understood that this product has not been evaluated on all varieties under all possible growing conditions. The user should exercise a responsible and cautious attitude when using this product on a given variety and variety under the performance under the growing conditions. **NO WARRANTY OR GUARANTEE IS MADE, EITHER ORAL OR WRITTEN, BY THE MANUFACTURER OF THIS PRODUCT WITH THE USE OF THE PRODUCT WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND EXCEPT FOR THE LIMITED WARRANTY SET FORTH IN THE STATE OF CALIFORNIA AND THE STATE OF ILLINOIS.** The O.M. Scott & Sons Company or any other dealer for any and all states, provinces or territories may differ from the use or handling of this product shall be the purchaser's responsibility. The user is responsible for the quality of this product purchased. The Buyer and all users are deemed to have accepted the terms of this label when they use or apply this application in working together. It is the responsibility of the user to read the label and follow the directions of the O.M. Scott & Sons Company.

For Full Directions, The O.M. Scott & Sons Company, Mayfield, Ohio 44130.

Suggested Application Settings	
Apply to moist foliage at the DOUBLE RATE when environmental conditions favor disease activity.	
Apply to moist foliage at the DOUBLE RATE when environmental conditions favor disease activity.	
Application Rate	1.0 lb/1000 sq ft (0.225 kg/1000 sq m)
Application Interval	2 weeks
Application Method	Drop spreader
Application Time	Moist foliage
Application Temperature	50°F to 85°F (10°C to 29°C)
Application Wind Speed	0 to 10 mph (0 to 16 km/h)
Application Humidity	Relative humidity > 50%
Application Rain	0.1 inch (2.5 mm) or more
Application Soil Type	All soil types
Application Soil pH	5.0 to 7.0
Application Soil Fertility	Normal
Application Soil Moisture	Moist
Application Soil Temperature	50°F to 85°F (10°C to 29°C)
Application Soil Type	All soil types
Application Soil pH	5.0 to 7.0
Application Soil Fertility	Normal
Application Soil Moisture	Moist
Application Soil Temperature	50°F to 85°F (10°C to 29°C)