

11/21

538-159

1/8 2

8510

Fungicide X



NOT REVIEWED
In Accordance with EPA Notice 82-2,
Based-on Draft Labeling Dated 5/28/91

Fungicide X

Fungicide X

Fungicide X



• prevents and controls brown patch, leaf spot, red leaf spot and *Fusarium* patch/pink snow mold (Washington, Oregon and California only)



Net Weight 52.86 lbs (23.97 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION: causes eye irritation. Do not get in eyes, on skin or on clothing. In case of contact, immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed.

Environmental Hazards: Do not apply directly to water. Do not contaminate water when cleaning of equipment washers. Do not apply when weather conditions favor drift from areas treated. Apply the product only as specified on this label.

ACTIVE INGREDIENT
Iproacone 3-(2,5-Dichlorophenyl)-N-(1-methyl-1H-imidazol-2-yl)-1H-imidazole-5-carboxamide 1.30%
INERT INGREDIENTS 98.70%
Total 100.00%

EPA Reg. No. 538-159 EPA Est. No. 538-OH-1 & 44616-MO-01
Subscription used is first letter of run code on end of container
Product of USA
ProTurf Division
The O.M. Scott & Sons Company
Marysville, Ohio 43041
© 1991 The O.M. Scott & Sons Company. All rights reserved.

8510

8510

Recommended For Use By Professional Turfgrass Managers

2012



Fungicide X

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Prevents and controls the following turfgrass diseases:

- brown patch
- leaf spot (*Helminthosporium* spp.)
- red leaf spot
- Fusarium* patch (Wash., Ore. and Calif. only)

When to use

To prevent brown patch and red leaf spot, apply at the **NORMAL** or **HEAVY RATE** (1.0 to 1.5 oz a.i./M) to moist or dry foliage when disease activity is anticipated and repeat at 2-week intervals. If heavy disease pressures are encountered, apply at the **HEAVY RATE** to moist or dry foliage. When diseases are brought under control, revert back to bi-weekly **NORMAL RATE** applications.

To prevent *Fusarium* patch, apply at the **LIGHT** or **NORMAL RATE** (0.5 to 1.0 oz a.i./M) to moist or dry foliage when disease activity is anticipated. Repeat at 2 to 3-week intervals.

To prevent leaf spot, apply to moist or dry foliage at the **LIGHT RATE** (0.5 oz a.i./M) when disease activity is anticipated. Repeat if needed after 3 to 4 weeks.

To control brown patch and red leaf spot, apply to moist or dry foliage at the **HEAVY RATE** (1.5 oz a.i./M) as soon as disease activity is noticed. Repeat bi-weekly until disease is under control then revert back to **NORMAL** or **HEAVY RATE** applications bi-weekly.

To control *Fusarium* patch, apply to moist foliage at the **NORMAL RATE** (1.0 oz a.i./M) when disease activity is first noticed. Repeat as needed at this rate at 2- to 3-week intervals. When disease activity is under control, revert to **LIGHT RATE** (0.5 oz a.i./M) at 2- to 3-week intervals.

To control leaf spot, apply to moist foliage at the **LIGHT** to **NORMAL RATE** (0.5 to 1.0 oz a.i./M) when disease activity is first noticed. Repeat as needed at intervals not closer than 2 weeks.

How to use

For **LIGHT**, **NORMAL** or **HEAVY RATE** applications (0.5, 1.0 or 1.5 oz a.i./M), apply to moist or dry foliage. Applications need not be watered in. Delay mowing for 24 hours after application to prevent particle pickup by mowers.

Expected time of disease activity

SPRING leaf spot and *Fusarium* patch
SUMMER leaf spot and brown patch
FALL leaf spot and *Fusarium* patch
WINTER *Fusarium* patch

Allow 1 week between application of this product and other control products. This guideline may be relaxed where a severe disease attack requires immediate treatment.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding down top and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. **NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE.** The exclusive remedy of the user or Buyer, and the limit of liability of The O.M. Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O.M. Scott & Sons Company.

ProTurf Division
 The O.M. Scott & Sons Company
 Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

52.66 lb (23.97 kg) treats 22,000 ft² at the **LIGHT RATE**

52.66 lb (23.97 kg) treats 11,000 ft² at the **NORMAL RATE**

52.66 lb (23.97 kg) treats 7,330 ft² at the **HEAVY RATE**

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTING		
			LIGHT RATE	NORMAL RATE	HEAVY RATE
Scotts Drop Type	3 mph	overlap wheels	5	7	9
Scotts Rotaries					
R-8, R-8A (Cone 9)	3 mph	9 feet	1 1/2	4	1 1/2
4-6 y. Models (See below)	4.5 mph	18 feet	3 1/2 in.	5 in.	6 in.
Vico™	4.5 mph	18 feet	21	27	34

Light Rate: 0.5 oz a.i./M, Normal Rate: 1.0 oz a.i./M, Heavy Rate: 1.5 oz a.i./M

Fungicide X

8510