


P.M. 31

538-159

128



ProTurf Fungicide VI XXXX

FRONT:

12/05/88

10/09/90 CR

05/04/91 JB/CR

(Scotts(R))
(ProTurf(R))
BRAND

Fungicide VI

o prevents and controls turfgrass diseases including benzimidazole tolerant dollar spot

NET WEIGHT 23.00 lbs (10.44 kg)
NET WEIGHT 52.86 lbs (23.97 kg) (optional bag size)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed.

Environmental Hazards: Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

ACTIVE INGREDIENT:

Iprodione 3-(3,5-Dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide.	1.30%
INERT INGREDIENTS:	98.70%
Total	100.00%

EPA Reg No 538-159

EPA Est 538-OH-1

Product of USA

ProTurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041

(c) 1991, The O.M. Scott & Sons Company, Marysville, Ohio 43041
All Rights Reserved.

Recommended for Use by Professional Turfgrass Managers

XXXX

Fungicide VI

Optional Front Panel Claims

- o prevents and controls brown patch, leaf spot, red leaf spot and Fusarium patch/pink snow mold (Washington, Oregon and California only)

OR

- o prevents and controls brown patch, leaf spot, red leaf spot, pink snow mold and gray snow mold

3, f8

RIGHT GUSSET:

XXXX

(Scotts(R))
(ProTurf(R))
BRAND

Fungicide VI

LEFT GUSSET:

(Scotts(R))
(ProTurf(R))
BRAND

Fungicide VI

XXXX

478

XXXX

Fungicide VI

(Scotts(R)
(ProTurf(R))
BRAND

Fungicide VI

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Prevents and controls the following turfgrass diseases:

brown patch
dollar spot (including
benzimidazole tolerant
strains)

leaf spot (Helminthosporium spp.)
red leaf spot
Fusarium patch (Wash, Ore, and
Calif Only)

DOLLAR SPOT, BROWN PATCH and RED LEAF SPOT: To prevent disease, apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

For active disease apply, to moist foliage at the DOUBLE RATE and lightly water in immediately to remove some of the product particles from the foliage. Repeat the application every 2 weeks. Once the disease is brought under control, use RECOMMENDED RATE applications at 2 week intervals for the remainder of the disease period.

FUSARIUM PATCH (Wash, Ore, and Cali only): To prevent disease, apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 to 3 week intervals throughout the expected disease period.

For active Fusarium patch, apply to moist foliage at the DOUBLE RATE and repeat at 2 to 3 week intervals. When the disease activity is brought under control, apply at the RECOMMENDED RATE at 2 to 3 week intervals.

LEAF SPOT (Helminthosporium spp.): To prevent disease, apply to moist foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat if needed after 3-4 weeks.

For active leaf spot, apply to moist foliage at the DOUBLE RATE when disease activity is first noticed. Repeat as needed at the DOUBE RATE at intervals not closer than 2 weeks.

REMINDER

Make all applications to moist foliage.

578
ALL DOUBLE RATE applications on putting green turf, except for the Fusarium patch and leaf spot, should be lightly watered in immediately after application. Heavy watering may significantly reduce disease control.

Delay mowing for 24 hours after application. This product is to be applied with drop spreaders only.

Do not graze treated area. Do not feed clippings to livestock. Keep children and pets off treated area until material has been washed off the foliage and the area is completely dry.

Please pay special attention to the handling information as noted in the warning statement to ensure safe use of this product.

Revised Optional Directions for Use

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This fungicide product prevents and controls the following turfgrass diseases:

- brown patch
- leaf spot
- red leaf spot
- pink snow mold (Fusarium nivale)
- gray snow mold (Typhula spp.)

For use on fairways, putting greens and other turf areas consisting of Kentucky bluegrass, bentgrass, perennial ryegrass, fine fescue and bermudagrass, or where these mixtures predominate. Product may also be applied to St. Augustinegrass turf areas.

When to Use

To prevent brown patch and red leaf spot, apply at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/M) to moist or dry foliage when disease activity is anticipated and repeat at 2-week intervals. If heavy disease pressures are encountered, apply at the HEAVY RATE to moist or dry foliage. When diseases are brought under control, revert back to bi-weekly NORMAL RATE applications.

To prevent pink and gray snow mold, apply at the LIGHT or NORMAL RATE (0.5 to 1.0 oz ai/M) to moist or dry foliage when disease activity is anticipated. Repeat at 2 to 3 week intervals.

For suppression of pink and gray snow mold, apply to dry foliage at the NORMAL or HEAVY RATE (1.0 to 1.5 oz. ai/M) prior to the first snow. Repeat in mid-winter when turf is free of snow and again immediately after the final snow melts in the spring.

Not a season long (all winter) control for snow molds unless applied on a 2 to 3 week application program.

To prevent leaf spot, apply to moist or dry foliage at the LIGHT RATE (0.5 oz ai/M) when disease activity is anticipated. Repeat if needed after 3 to 4 weeks.

To control brown patch and red leaf spot, apply to moist or dry foliage at the HEAVY RATE (1.5 oz ai/M) as soon as disease activity is noticed. Repeat bi-weekly until disease is under control, then revert back to NORMAL or HEAVY RATE applications bi-weekly.

To control pink and gray snow mold, apply to dry foliage at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/M) when disease activity is first noticed. Repeat as needed at this rate at 2 to 3 week intervals. When disease activity is under control, revert to the LIGHT or NORMAL RATE (0.5 to 1.0 oz ai/M) at 2 to 3 week intervals.

To control leaf spot, apply to moist foliage at the NORMAL to HEAVY RATE (1.0 to 1.5 oz ai/M) when disease activity is first noticed. Repeat as needed at intervals not closer than 2 weeks.

How to Use

LIGHT, NORMAL or HEAVY RATE applications (0.5, 1.0 or 1.5 oz ai/M), apply to moist or dry foliage for summer diseases. Applications need not be watered in. Apply to dry foliage for snow mold prevention/control.

Delay mowing for 24 hours after application to prevent particle pickup by mowers.

Expected time of disease activity:

Spring -- leaf spot, pink snow mold (Fusarium patch), gray snow mold

Summer -- leaf spot and brown patch

Fall -- leaf spot and pink snow mold (Fusarium patch)

Winter -- pink snow mold (Fusarium patch), and gray snow mold

Allow one week between the application of this and other control products. This guideline may be relaxed where a severe disease attack requires immediate treatment.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

528
1

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide the necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IF MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O.M. Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O.M. Scott & Sons Company.

ProTurf Division
The O.M. Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application

23 lbs (10.44 kg) treats 11,000 ft² (1/4 acre/1022 m²) at RECOMMENDED RATE
23 lbs (10.44 kg) treats 5,500 ft² (1/8 acre/ 511 m²) at DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			RECOMMENDED	DOUBLE
Scotts Drop	3 mph	overlap wheels	6 3/4	9 1/2

(Optional Bag Size and Spreader Settings)

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application

52.86 lbs (23.97 kg) treats 22,000 ft² at the LIGHT RATE
52.86 lbs (23.97 kg) treats 11,000 ft² at the NORMAL RATE
52.86 lbs (23.97 kg) treats 7,330 ft² at the HEAVY RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			LIGHT RATE	NORMAL RATE	HEAVY RATE
Scotts Drop Type	X	X	X	X	X
Scotts Rotaries	X	X	X	X	X
Lely Models (see below)	X	X	X	X	X

Lely Models WTR, WFR, HR and 1250. PTO at 450 RPM