

Pm 21

538-159

ProTurf Fungicide VI XXXX

1 of 8

FRONT:

**ACCEPTED**  
 DEC 4 1990  
 538-159

12/05/88

10/09/90 CR

(Scotts(R))  
(ProTurf(R))  
BRAND

Fungicide VI

o prevents and controls turfgrass diseases including benzimidazole tolerant dollar spot

NET WEIGHT 23.00 lbs (10.44 kg)  
NET WEIGHT 52.86 lbs (23.97 kg) (optional bag size)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed.

Environmental Hazards: Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

ACTIVE INGREDIENT:

Iprodione 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-	
2,4-dioxo-1-imidazolidinecarboxamide. . . . .	1.30%
INERT INGREDIENTS: . . . . .	98.70%
Total	100.00%

EPA Reg No 538-159

EPA Est 538-OH-1

Product of USA

ProTurf Division  
The O. M. Scott & Sons Company  
Marysville, Ohio 43041

(c) 1990, The O.M. Scott & Sons Company, Marysville, Ohio 43041  
All Rights Reserved.

Recommended for Use by Professional Turfgrass Managers

XXXX

Fungicide VI

Optional Front Panel Claims

- prevents and controls brown patch, leaf spot, red leaf spot and Fusarium patch/pink snow mold (Washington, Oregon and California only)

308

RIGHT GUSSET:

XXXX

(Scotts(R))  
(ProTurf(R))  
BRAND

Fungicide VI

LEFT GUSSET:

(Scotts(R))  
(ProTurf(R))  
BRAND

Fungicide VI

XXXX

XXXX

Fungicide VI

(Scotts<sup>(R)</sup>)  
(ProTurf<sup>(R)</sup>)  
BRAND

Fungicide VI

Directions for use

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Prevents and controls the following turfgrass diseases:

brown patch  
dollar spot (including  
benzimidazole tolerant  
strains)

leaf spot (Helminthosporium spp.)  
red leaf spot  
Fusarium patch (Wash, Ore, and  
Calif only)

DOLLAR SPOT, BROWN PATCH and RED LEAF SPOT: To prevent disease, apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

For active disease apply, to moist foliage at the DOUBLE RATE and lightly water in immediately to remove some of the product particles from the foliage. Repeat the application every 2 weeks. Once the disease is brought under control, use RECOMMENDED RATE applications at 2 week intervals for the remainder of the disease period.

FUSARIUM PATCH (Wash, Ore, and Cali only): To prevent disease, apply to moist foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 to 3 week intervals throughout the expected disease period.

For active Fusarium patch, apply to moist foliage at the DOUBLE RATE and repeat at 2 to 3 week intervals. When the disease activity is brought under control, apply at the RECOMMENDED RATE at 2 to 3 week intervals.

LEAF SPOT (Helminthosporium spp.): To prevent disease, apply to moist foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat if needed after 3-4 weeks.

For active leaf spot, apply to moist foliage at the DOUBLE RATE when disease activity is first noticed. Repeat as needed at the DOUBE RATE at intervals not closer than 2 weeks.

REMINDER

Make all applications to moist foliage.

538  
ALL DOUBLE RATE applications on putting green turf, except for the Fusarium patch and leaf spot, should be lightly watered in immediately after application. Heavy watering may significantly reduce disease control.

Delay mowing for 24 hours after application. This product is to be applied with drop spreaders only.

Do not graze treated area. Do not feed clippings to livestock. Keep children and pets off treated area until material has been washed off the foliage and the area is completely dry.

Please pay special attention to the handling information as noted in the warning statement to ensure safe use of this product.

## Optional Directions for Use

### When to Use

To prevent brown patch and red leaf spot, apply at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/M) to moist or dry foliage when disease activity is anticipated and repeat at 2-week intervals. If heavy disease pressures are encountered, apply at the HEAVY RATE to moist or dry foliage. When diseases are brought under control, revert back to bi-weekly NORMAL RATE applications.

To prevent Fusarium patch, apply at the LIGHT or NORMAL RATE (0.5 to 1.0 oz ai/M) to moist or dry foliage when disease activity is anticipated. Repeat at 2 to 3 week intervals.

To prevent leaf spot, apply to moist or dry foliage at the LIGHT RATE (0.5 oz ai/M) when disease activity is anticipated. Repeat if needed after 3 to 4 weeks.

To control brown patch and red leaf spot, apply to moist or dry foliage at the HEAVY RATE (1.5 oz ai/M) as soon as disease activity is noticed. Repeat bi-weekly until disease is under control, then revert back to NORMAL or HEAVY RATE applications bi-weekly.

To control Fusarium patch, apply to moist foliage at the NORMAL RATE (1.0 oz ai/M) when disease activity is first noticed. Repeat as needed at this rate at 2 to 3 week intervals. When disease activity is under control, revert to LIGHT RATE (0.5 oz ai/M) at 2 to 3 week intervals.

To control leaf spot, apply to moist foliage at the LIGHT to NORMAL RATE (0.5 to 1.0 oz ai/M) when disease activity is first noticed. Repeat as needed at intervals not closer than 2 weeks.

### How to Use

For LIGHT, NORMAL or HEAVY RATE applications (0.5, 1.0 or 1.5 oz ai/M), apply to moist or dry foliage. Applications need not be watered in.

Delay mowing for 24 hours after application to prevent particle pickup by mowers.

Expected time of disease activity:

Spring -- leaf spot and Fusarium patch

Summer -- leaf spot and brown patch

Fall -- leaf spot and Fusarium patch

Winter -- Fusarium patch

Allow one week between the application of this and other control products. This guideline may be relaxed where a severe disease attack requires immediate treatment.

748

## Storage and disposal

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division      The O.M. Scott & Sons Company      Marysville, Ohio 43041

---

## IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide the necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IF MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O.M. Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O.M. Scott & Sons Company.

ProTurf Division  
The O.M. Scott & Sons Company  
Marysville, Ohio 43041

---

8.14

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application

23 lbs (10.44 kg) treats 11,000 ft<sup>2</sup> (1/4 acre/1022 m<sup>2</sup>) at RECOMMENDED RATE  
 23 lbs (10.44 kg) treats 5,500 ft<sup>2</sup> (1/8 acre/ 511 m<sup>2</sup>) at DOUBLE RATE

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>	
			<u>RECOMMENDED</u>	<u>DOUBLE</u>
Scotts Drop	3 mph	overlap wheels	6 3/4	9 1/2

**(Optional Bag Size and Spreader Settings)**

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application

52.86 lbs (23.97 kg) treats 22,000 ft<sup>2</sup> at the LIGHT RATE  
 52.86 lbs (23.97 kg) treats 11,000 ft<sup>2</sup> at the NORMAL RATE  
 52.86 lbs (23.97 kg) treats 7,330 ft<sup>2</sup> at the HEAVY RATE

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>		
			<u>LIGHT RATE</u>	<u>NORMAL RATE</u>	<u>HEAVY RATE</u>
Scotts Drop Type	X	X	X	X	X
Scotts Rotaries	X	X	X	X	X
Lely Models (see below)	X	X	X	X	X

Lely Models WTR, WFR, HR and 1250. PTO at 450 RPM