

PROTURF FUNGICIDE 7 #8645
02/04/85 MB 04/28/86 JF
01/23/86 JF
04/21/86 JF

FRONT:

(Scotts (R))
ProTurf (R)
BRAND

PM 71
53--161
p. 1. of 5

Fungicide 7

systemic fungicide that prevents and controls dollar spot (including benzimidazole tolerant strains), brown patch, anthracnose and rust

Net Weight 29 3/16 lbs (13.24 kg)

LEFT COLUMN

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. If in eyes, flush with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated area. Do not feed clippings to livestock. Keep children and pets off treated area until material has been washed off the foliage and the area is completely dry.

Environmental Hazards: Do not apply directly to water. Do not contaminate water by cleaning of equipment or

RIGHT COLUMN

disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

ACCEPT

JAN 13 1987

U.S. Environmental Protection Agency
Registration Division
Washington, D.C. 20460
or call 1-800-455-6114 for the nearest
registered user.

BEST AVAILABLE COPY

FRONT CONTINUED:

pg 2 of 5

ACTIVE INGREDIENT:

(1-(Chlorophenoxy)-3,3-dimethyl-1-
(1H-1,2,4-triazol-1-yl)-2-butanone

.59%

INERT INGREDIENTS:

99.41%

Total 100.00%

EPA Reg No 538-161
Product of USA

EPA Est 538-OH-1

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

(c)1986. The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professionals

8645

Fungicide 7

GUSSETS:

RIGHT GUSSET

8645

(Scotts(R))
ProTurf(R)
BRAND

Fungicide 7

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

Fungicide 7

8645 (Packaging Calendar)

BACK:

Fungicide 7

8645

(Scotts(R))
ProTurf(R)
BRAND

Fungicide 7

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on turfgrass areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine fescue, bermudagrass or mixtures where these grasses predominate.

TO PREVENT DOLLAR SPOT, ANTHRACNOSE AND RUST apply to moist or dry foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT, ANTHRACNOSE AND RUST apply to moist or dry foliage at two week intervals. Use the DOUBLE RATE for light to moderate disease infestations and the HEAVY RATE for severe disease infestations. When the disease activity is under control, apply at the RECOMMENDED RATE at two week intervals.

TO PREVENT BROWN PATCH apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

RIGHT COLUMN

TO CONTROL ACTIVE BROWN PATCH apply to moist or dry foliage at the HEAVY RATE every two weeks. When the disease activity is under control, apply at the DOUBLE RATE at two week intervals.

To avoid product pick-up lightly irrigate treated areas soon after application.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding down top and securing. Protect from water and extreme heat.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

BEST AVAILABLE COPY

BACK CONTINUED:

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings
to provide proper distribution calibrate spreader before application .

29 3/16 lbs treats 22,000 ft² (1/2 acre) at the RECOMMENDED RATE
29 3/16 lbs treats 11,000 ft² (1/4 acre) at the DOUBLE RATE
29 3/16 lbs treats 5,500 ft² (1/8 acre) at the HEAVY RATE

<u>SPREADER</u>	<u>GROUND SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>		
			<u>RECOMMENDED</u>	<u>DOUBLE</u>	<u>HEAVY</u>
Scotts Drop Scotts Rotaries	3 mph	overlap wheels	4 1/4	5 3/4	7 1/4
R-7, R-7X	3 mph	9 ft	D	G	M
R-8 (cone 8)	3 mph	9 ft	J	L	O 1/2
Lely Models (see below)	4.5 mph	16 ft	3 1/4 IV	4 1/4 IV	5 3/4 IV

Lely Models WTR, WFR, HR and 1250. Run at 450 rpm.
Metric coverage: 13.24 kg treats 2044 m² at the RECOMMENDED RATE, 1022 m² at the DOUBLE RATE and 511 m² at the HEAVY RATE.