

ACCEPTED
 JUL 17 1981
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the reasons stated, registered under EPA Reg. No. 538-169

Scotts
spot grass & weed control

spot grass & weed control

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For easy control of undesirable weeds and grasses. Avoid application to foliage of desirable plants and vegetable gardens.

SHAKE WELL. Invert can to direct spray on leaves and crown of plants. Wet foliage thoroughly with a 2 or 3 second burst of spray. Avoid watering for 24 hours to insure effectiveness.

- Severe browning of treated vegetation occurs within 12 hours. In cool weather, action is slower.
- Repeat treatments in 5-7 days or as regrowth occurs. See side panel for treatments required.
- It is not necessary (or desirable) to remove dead weeds and grasses. They will decay and eventually disappear.
- Grass seed can be planted anytime after weeds or other plants are controlled.

LIMITATIONS:

- Do not apply on a windy day. Apply on a calm day to avoid drift or application on desirable lawn grasses or nearby plants.
- Grasses within the treated areas will brown and die.
- Do not track material from treated area to untreated area.
- Will not control weed seeds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals: Avoid breathing spray mist. Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep children and pets off treated area until after first rain. Watering is recommended after 48 hours if it has not rained.

Environmental Hazards: Keep out of lakes, streams, and ponds. Apply this product only as specified on the label.

Storage and Disposal: Do not reuse empty container. Wrap container and put in trash container.

Physical Hazards: Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

The O M Scott & Sons Company
 Marysville, Ohio 43041

Products of U S A

stops grass and weeds in walks, driveways, lawns



ACTIVE INGREDIENTS:		Total arsenic expressed as elemental in all water soluble form 0.71%
Sodium cacodylate	1.26%	EPA Reg. No. 538-169
Dimethylarsinic acid	21%	EPA Est. XXXX XX X
NET INGREDIENTS	98.53%	Distributed by:
Total	100.00%	The O M Scott & Sons Company
		Marysville, Ohio 43041

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
 See other side for precautions

NET WT.
1 LB. 4 OZ.
(.567 kg.)

One Treatment

- annual bromegrass
- barnyardgrass
- brassbuttons • buckhorn
- chickweed (common, mouseear)
- chicory
- crabgrass (hairy, smooth)
- dandelion • dichondra
- dog fennel • dollarweed
- English daisy • fall panicum
- fleabane • Florida pusley
- foxtail • goosegrass
- knotweed • lambsquarters
- lovegrass
- pigweed
- plantain • poa annua
- purslane • fine fescues
- smartweed • speedwell
- spotted spurge
- stinging nettle
- wild carrot • witchgrass
- yellow woodsorrel (oxalis)

Two or More Treatments:

- bahiagrass • bentgrass
- bermudagrass • clover
- coarse fescues • dallisgrass
- false dandelion
- Kentucky bluegrass
- kikuyugrass • nimblewill
- nutsedge (yellow)
- quackgrass • ryegrass
- St. Augustinegrass
- thistle (Canada) • timothy
- velvetgrass • wild onion

Also use Scotts' Spot Dandelion Control for selective control of certain undesirable lawn weeds.

Scotts Promise to you—satisfaction or your money back.

© 1981, The O M Scott & Sons Company
 All rights reserved 6515