

XXXX



ACCEPTED
MAY 7 1982
Insecticide
Labeling
Act



Fertilizer
Plus
Fungicide 7

Fertilizer Plus Fungicide 7

Fertilizer
Plus
Fungicide 7



- provides systemic disease prevention and control
- provides controlled release nitrogen feeding

Net Weight 27½ lbs
(12.47 kg)

XXXX

XXXX

For Professional Use Only

XXXX

XXXX



Fertilizer Plus Fungicide 7

Directions for use

Scatter or broadcast fertilizer and fungicide product in a uniform, consistent manner over the lawn.

For use on lawns, trees, tees, and shrubs consisting of Kentucky bluegrass, annual bluegrass, fine fescue, perennial ryegrass, tall fescue, and doggrass, or mixtures. Avoid use on these grasses in pastures.

TO PREVENT DOLLAR SPOT, ANTHRACNOSE AND RUST (Puccinia spp.)

Apply at the **RECOMMENDED RATE** when environmental conditions favor disease activity. Repeat at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT, ANTHRACNOSE AND RUST (Puccinia spp.)

Apply at the **DOUBLE RATE** to moist or dry foliage at 2 week intervals. After the disease activity subsides, return to the **RECOMMENDED RATE** at two week intervals.

TO PREVENT BROWN PATCH

Apply at the **DOUBLE RATE** to moist or dry foliage when environmental conditions favor disease activity. Repeat at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE BROWN PATCH

Use a non-fertilizer fungicide such as ProTurf Fungicide 7. Use the amount of fertilizer application. High levels of nitrogen fertilizer may increase the disease risk.

Do not use on lawns that are being treated with a fungicide.

RECOMMENDED RATE

Apply at the rate of 1.5 lbs. per 1,000 sq. ft. of lawn area. For broadcast application, use a broadcast spreader. For spot application, use a hand-held spreader. Do not apply to dry lawns when temperature is above 90°F.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

SPRAYS: Store in a cool, dry place. Release vapors by fanning, blowing, and stirring. **Protect from water and extreme heat.**


WATER CONTAMINATION: Do not use if these waters are used for drinking. Do not use if these waters are used for irrigation. Do not use if these waters are used for watering lawns.

WATER CONTAMINATION: Do not use if these waters are used for drinking. Do not use if these waters are used for irrigation. Do not use if these waters are used for watering lawns.

WATER CONTAMINATION: Do not use if these waters are used for drinking. Do not use if these waters are used for irrigation. Do not use if these waters are used for watering lawns.

WATER CONTAMINATION: Do not use if these waters are used for drinking. Do not use if these waters are used for irrigation. Do not use if these waters are used for watering lawns.

Suggested sprayer settings



1.5 lbs. per 1,000 sq. ft.

1.5 lbs. per 1,000 sq. ft.

XXXX