

ACCEPTED

OCT 25 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act
EPA Reg. No. 280.1274



28-0-12
Fertilizer Plus
Fungicide 7

8307



28-0-12
Fertilizer Plus
Fungicide 7

High Density

Scotts ProLurf
28-0-12
Fertilizer Plus
Fungicide 7

28-0-12 Fertilizer Plus Fungicide 7

28-0-12
Fertilizer Plus
Fungicide 7



- provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot
- provides controlled release nitrogen feeding

Net Weight 27 1/2 lbs (12.47 kg)

8307

8307

For Professional Use Only

8307

28-0-12
Fertilizer Plus
Fungicide 7

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on turf areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine leaf fescue, bermudagrass or mixtures where these grasses predominate.

TO PREVENT DOLLAR SPOT ANTHRACNOSE AND RUST: Purchase and apply to moist or dry foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT ANTHRACNOSE AND RUST: Purchase and apply to moist or dry foliage at the DOUBLE RATE to moist or dry foliage at two week intervals. When the disease activity is under control, apply at the RECOMMENDED RATE at two week intervals.

TO PREVENT BROWN PATCH: apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE BROWN PATCH: Use a non-fertilizer fungicide such as ProLurf Fungicide 7 to limit the amount of nitrogen applied to turf during or just prior to the onset of hot, humid weather may increase the severity of

brown patch. Follow with the RECOMMENDED RATE of either product once the disease is controlled.

Note carefully

To avoid product pick-up, lightly irrigate treated areas soon after application. On short cut turfgrass (less than 1/2"), apply only to dry foliage when temperatures are above 60°F.

Storage and disposal

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reveal opened bags by folding down top and securing. PROTECT FROM WATER AND EXTREME HEAT.

PESTICIDE DISPOSAL: Pesticide or mixed waste that cannot be used, according to label instructions, must be disposed of according to applicable federal, state or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into appropriate equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

ProLurf Division
The Scotts Company
Marysville, Ohio 44041

Suggested Sprayer Settings			
Application Method	Rate (lb/1000 sq ft)	Pressure (psi)	Flow (gpm)
Hand Sprayer	1.0	10-15	0.5
Knapsack	1.0	10-15	0.5
Backpack	1.0	10-15	0.5
Tractor Sprayer	1.0	10-15	0.5
Tractor Sprayer	1.0	10-15	0.5
Tractor Sprayer	1.0	10-15	0.5

BEST DOCUMENT AVAILABLE