Please read instructions on		38-775 Dilating form	6/30/98 Form Appr	ved OMB No 2		1 POPE 1 7- 2 pproval expires 05-31-98
	Unite Environmental	ed States		Registra Amend Cother	ation	OPP Identifier Number
		Application f	or Pesticide - Sectio	n l		
1. Company/Product Number 538- 175			2. EPA Product Manager 3. Proposed Classification Joanne Miller		posed Classification	
4. Company/Product (Name) Super Plus 2 For Grass			РМ# РМ-23			
<ol> <li>Name and Address of Applicant (Include ZIP Code)</li> <li>The Scotts Company</li> <li>14111 Scottslawn Road</li> <li>Marysville, OH 43041</li> </ol>			6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name			
Check if this	is a new address		Section II		·	
Amendment - Explain below.		Final printed lab	Final printed labels in response <b>NOTIFICATION</b> XX-XX-XX "Me Too" Application JUN 3 0 1998 Other - Explain below.			
	additional fert	ilizer ratio to	ecord: "26-4-4 Ferti the record: "26-4-4" Section III			
Child-Resistant Packaging Yes* No *Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per	Water Soluble Packaging Yes No If "Yes" No. pe Package wgt. Contai		e of Containe Metal Plastic Glass Paper	
Location of Net Contents In	nformation Container	4. Size(s) Reta 50.0 lb, and		5. Location o On Lat	bel	
6. Manner in Which Label is /	Affixed to Product	Lithograph				
·			Section IV			
1. Contact Point <i>(Complete items directly below for identificatio</i> Name T Chris J. Wibłe		Title			ss this application.) Telephone Nc. (Include Area Code) (907) €14-7214	
acknowledge that any kr under applicable law.		ading statement ma	on Ihments thereto are true, accu ay be punishable by fine or im Title Senior Regulato	prisonment or bot		6. Dete Application Received (Stam))ed)
4. Typed Name	J. Wible	5.	Date June 22,	1998		



### NOTIFICATION

2 73

JUN 3 0 1998

### 26-4-4 Fertilizer Plus Weed Control

### KEEPS LAWNS GREENER LONGER

\*Compared to fertilizers without slow-release nitrogen

DO NOT USE ON St. Augustinegrass, dichondra or bentgrass lawns.

### **ACTIVE INGREDIENTS:**

EPA Reg. No.538-175 EPA Est. 55640-OH-002

# KEEP OUT OF REACH OF CHILDREN

See back panel for additional precautionary statements

KILLS DANDELIONS AND 50 OTHER TYPES OF WEEDS 50.0 lb Treats 13,360 sq.ft.

26-4-4

)

### **GUARANTEED ANALYSIS**

Total nitrogen (N)	
5.0% other water soluble nitrogen**	
0.4% water insoluble nitrogen	
Available phosphate (P2O5) 4%	
Soluble potash (K2O) 4%	
Sulfur (S)	
5.0% free sulfur (S)	

Derived from monoammonium phosphate, urea, polymer-coated sulfur-coated urea, methylene ureas, and muriate of potash. \*A portion of the urea has been coated to provide 7.7% coated-

slow release nitrogen. \*\*Contains 4.1% slowly available methylenediurea and dimethylenetriurea nitrogen.

U. S. Pat. Nos. 4,378,238	5,219,465	5,102,440
5,300,135	5,405,426	

Distributed by: The Scotts Company 14111 Scottslawn Rd. Marysville, OH 43041 APPLICATION RATE per 1,000 sq. ft. (20'X50') For established lawns-3.74 lbs (15 cups) per 1,000 sq. ft.

### Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds of 2, 4-D per acre per application. The maximum number of broadcast applications per treatment site is two per year.

- Apply when weeds are actively growing but not before yellow dandelion blooms have appeared.
- Apply to moist or dry foliage, however, for best results, apply to moist foliage.
- Be careful when applying this product near shrubs, flower beds or vegetable gardens as weed control may damage these desirable plants.
- Do not over apply, especially under the drip zone of desirable shrubs and trees as some of the control ingredients are root absorbed.
- Don't use on dichondra, St. Augustine, or bentgrass lawns. Don't use on newly seeded or newly sodded lawns until after the fourth mowing.
- · Don't plant grass for one month after using this product.
- Don't apply when daytime temperatures are consistently 90 degrees or above.
- Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product. This product should not be applied if rain is expected within the next 24 hours.
- · Allow at least 4-6 weeks between program applications.

#### Storage and Disposal

Storage: Store in a clean, dry place. Reseal opened container or bag by folding top down and securing.

<u>Pesticide Disposal:</u> Securely wrap original container in several layers of newspaper and discard in trash.

. . . . .

· , , , , ,

<u>Container Disposal:</u> Do not reuse bag. Discard bag in trash.

## Net Wt. 50.0 lb (22.6 Kg)



### 26-4-4 Fertilizer Plus Weed Control

# 3 IMPORTANT REASONS TO USE 26-4-4 FERTILIZER PLUS WEED CONTROL

1. Clears out dandelions and more than 50 broadleaf weeds.

Controls more than 50 common lawn weeds such as dandelion, clover and chickweed. Effective on both large and small weeds. (See list below.) Kills roots and all.

### 2. Lawns stay greener longer.

Provides extended feeding for extra long lasting <u>nitro-</u> <u>gen</u> greening. One feeding lasts for months. Saves work, saves money.

### 3. No burning. No surge growth.

Each fertilizer particle provides safe gradual release that won't burn or cause surge growth when applied as directed.

### **RESULTS YOU CAN SEE**

Your lawn will be greener in 4 to 6 days. Dandelions and other weeds will be killed in 3 to 4 weeks.

#### HOW WEED CONTROL WORKS

Small, lightweight granules mean better weed control. Most lawn weed controls must be absorbed through the weed's leaves to be effective. The weed control in this product consists of granules that are small and lightweight, so they will stay on even the smallest weed leaves. The weed control is absorbed by the weed's leaves, killing the plant completely, root and all.

### WHAT WEEDS ARE CONTROLLED

Controls more than 50 broadleaf weeds:

aster bindweed bittercress black medic bristly oxtongue buckhorn bull thistle burclover buttonweed Canada thistle carpetweed catsear (coast dandelion) centella chickweed chicory cinquefoil clover (hop) clover (red) clover (white) cudweed curly dock dandelion

)

dichondra prickly lettuce dollarweed puncturevine English daisy purslane eveningprimrose pusley (Florida) filaree pussytoes fleabane ragweed around ivy red sorrel hawkweed (sheep sorrel) scarlet pimpernel healall henbit shadflower (whitlowort) horseweed knotweed(seedling) shepherdspurse smartweed lambsquarter lespedeza(seedling) sowthistle lippia(matchweed) spurweed mallow stitchwort threeseeded moneywort mercury morning glory mustard thymeleaved pennywort sandwort pepperweed vetch pigweed wild carrot plantain varrow vellow rocket

### When Using A Drop Spreader:

Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Then operate spreader the longest way of the lawn. Overlap wheel marks to insure uniform feeding. Shut off spreader when you have reached the header strips or when you have stopped.

3

8

, Э

#### When Using A Rotary Spreader:

Shut-off spreader any time you stop. For complete coverage make each pass according to spreader manufacturer's directions so some overlapping occurs.

### PRECAUTIONARY STATEMENTS

<u>Hazards to Humans and Domestic Animals</u>: CAUTION. Causes eye injury. Avoid contact with eyes, skin or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs.

Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with scap and water.

Environmental Hazards: Do not apply when weather conditions favor drift from target area. Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

SETTINGS FOR DROP SPREADERS	NORMAL
Scotts - all models	
SETTINGS FOR ROTARY SPREADERS	NORMAL
Scotts EasyGreen "(30" width)	4 1/2

### CAN WE HELP YOU? Call toll-free 1-800-543-TURF

Proudly produced in the U. S. A. ©1998, The Scotts Company. World Rights Reserved.

. . . .

,

ı –

y na sina

. .

June 18, 1998