

FRONT:

FLUID FUNGICIDE 83386
07/16/84 RF 12/17/84 MJB
08/01/84 RF 03/25/85 JF
08/06/84 RF 07/28/86 JF

(Scotts(R))
ProTurf(R)
BRAND

83386

Fluid Fungicide

prevents and controls turf diseases on tees, greens, fairways and similar turf areas

Net Contents 1/2 Gallon (1.89 liters)

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Do not get in eyes. If in eyes, flush with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed.

Environmental Hazards: Do not apply directly to water.

ACTIVE INGREDIENTS:

Thiophanate-methyl, Dimethyl [(1,2-phenylene) bis (iminocarbonothioyl)] bis[carbamate]	19.65%
Iprodione, 3-(3,5-Dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide	19.65%
INERT INGREDIENTS:	60.70%
Total	100.00%

EPA Reg No 538-183


EPA Est 37429-GA-1

Product of USA

Sold by: ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

(c)1984. The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professionals



ACCEPTED
FEB 1 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 538-183

BACK:

Directions for use

Read entire label and use only as directed

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Prevents and controls the following diseases on the turfs where they occur:

leaf spot/melting out	(<u>Helminthosporium spp./Drechslera spp.</u>)
dollar spot	including benzimidazole tolerant strains
brown patch	
pink snow mold	(Washington, Oregon, California only)
anthracnose	(on <i>Poa annua</i>)

Mixing instructions

Shake product container well before pouring into spray tank.

Mix ProTurf Fluid Fungicide into the spray tank during filling, while maintaining agitation.

Use with 4-7 gallons of water per 1000 ft² for tees and greens. Water requirement can be reduced to 2 gallons per 1000 ft² for larger turf areas.

Thoroughly rinse container and pour remaining contents into spray tank.

Do not combine ProTurf Fluid Fungicide in the spray tank with surfactants or other adjuvants.

Application directions


Apply with standard pressurized spray equipment.

To Prevent Disease - Treat turf at the normal rate when weather conditions are favorable for the development of disease. Repeat on a 10-14 day schedule.

To Control Mild Disease Infestations - Use the medium rate. Repeat the medium rate on a 10-14 day schedule until the disease is brought under control, then treat at the normal or medium rate on a preventive schedule.

To Control Moderate to Severe Disease Infestations - Use the heavy rate. Repeat on a 7-10 day schedule until disease is brought under control, then treat at the normal or medium rate on a preventive schedule.

Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.



BACK CONTINUED:

Coverage Chart for the Three Recommended Rates of ProTurf Fluid Fungicide

Rate (oz./M)	1 Pint (16 fl. oz.) Will Cover	1 Quart (32 fl. oz.) Will Cover	2 quarts (64 fl. oz.) Will Cover
Normal (1.1 fl. oz./M)	15,000 ft ²	30,000 ft ²	60,000 ft ²
Medium (1.6 fl. oz./M)	10,000 ft ²	20,000 ft ²	40,000 ft ²
Heavy (2.1 fl. oz./M)	7,500 ft ²	15,000 ft ²	30,000 ft ²

Use with 4-7 gallons of water per 1000 ft² for tees and greens. Water requirement can be reduced to 2 gallons per 1000 ft² for larger turf areas.

Limitations: Not a control for Pythium, stripe smut, spring dead spot, gray snow mold or Fusarium blight.

Note to User


Do not allow spray mixture to stand in the tank for longer than 12 hours.

Spray mixture should be agitated continuously during application.

Use a properly calibrated sprayer.

For best results do not water or mow treated areas within 24 hours of application.

Except in cases of severe infestation, allow one week between applications of this and other control products.



BACK CONTINUED:

Storage and disposal

STORAGE: Store in a cool, clean, dry place. Do not store at temperatures exceeding 120°F nor at temperatures below 32°F.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state or local authorities, by burning. If burned, stay out of smoke. Do not reuse empty container.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

83386

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide the necessary data to support its use as described on the label. However, the user should understand that tests have not been carried out under all possible use conditions, and that Scotts cannot and does not warrant that the product is compatible with all other chemicals. User should conduct its own compatibility testing and should test its mixes and equipment thoroughly before using the product on customers' property. ACCORDINGLY, SCOTTS MAKES NO WARRANTY, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH THESE DIRECTIONS AND THE DIRECTIONS ON THE LABEL. The exclusive remedy of the user and buyer and limit of liability of The O M Scott & Sons Company, or any other seller, for any and all losses, damages, and injuries resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by an officer of The O M Scott & Sons Company.

BEST AVAILABLE COPY