

FRONT CONTINUED:

52.8-122

Guaranteed Analysis

Total nitrogen (N) . . . . . 32%  
0.7% ammoniacal nitrogen  
21.3% urea, methylene ureas nitrogen  
10.0% water insoluble nitrogen  
Available phosphoric acid (P<sub>2</sub>O<sub>5</sub>) . . . . . 4%  
Soluble potash (K<sub>2</sub>O) . . . . . 4%

Derived from: monoammonium phosphate, urea, methylene ureas, and potassium sulfate.

Product of USA.  
EPA Est 538-OH-1

EPA Reg. No. 538-XXX

ProTurf Division      The O M Scott & Sons Company      Marysville, Ohio 43041

(c) 1984      The O M Scott & Sons Company      All rights reserved.

For Professional Use Only

XXXX

GUSSETS

RIGHT GUSSET:

(Scotts (R))  
ProTurf  
Brand

32-4-4  
Fertilizer Plus  
Dicot Weed Control III

XXXX

LEFT GUSSET:

(Scotts (R))  
ProTurf  
Brand

32-4-4  
Fertilizer Plus  
Dicot Weed Control III

XXXX

BACK:

32-4-4

Fertilizer Plus  
Dicot Weed Control III

XXXX

(Scotts (R))  
ProTurf  
Brand

32-4-4  
Fertilizer Plus  
Dicot Weed  
Control III

---

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on golf course putting greens. Fertilizes grass and controls these rosette and vining type weeds:

Aster	English daisy	Pigweed
Bittercress	Eveningprimrose	Plantain
Black medic	Filaree	Prickly lettuce
Bristly oxtongue	Fleabane	Purslane
Buckhorn	Ground ivy	Pusseytoes
Bull thistle	Hawkweed	Ragweed
Burclover	Healall	Scarlet pimpernel
Buttonweed	Henbit	Shadflower (whitlowort)
Carpetweed	Horseweed	Shepherdspurse
Catsear (coast dandelion)	Knotweed (seedling)	Smartweed
Chickweed	Lambsquarter	Sowthistle
Chickory	Lippia (matchweed)	Stitchwort
Cinquefoil	Mallow	Threeseeded mercury
Clover (white)	Moneywort	Thymeleaf sandwort
Cudweed	Morningglory	Vetch
Curly dock	Mustard	Yellow rocket
Dandelion	Pennywort	Yellow toadflax
Dichondra	Pepperweed	

For use on ornamental Turfgrass areas other than putting greens.

Apply to moist foliage at the RECOMMENDED RATE or for better distribution when using a rotary spreader, apply at the HALF RATE with a reduced width of coverage (see spreader settings). Apply when weeds are growing actively. Grass should not be so tall it touches the bottom of the spreader.

Rain, sprinkling, mowing, or heavy traffic within 24 hours after application can knock granules off weeds and may reduce effectiveness. Apply under non-windy conditions, thereby reducing drift to non-target areas, and optimizing particle distribution for good weed control. Retreating may be necessary, but wait at least 3 weeks to observe full results of first application.

BACK CONTINUED:

**PRECAUTIONS**

Avoid application when daytime temperatures are consistently 90° or above.

Do not over apply, especially under the branch spread of trees and shrubs.

On new grass plantings delay application until after the fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 1 month after application.

Not for use on creeping bentgrass, St. Augustinegrass, carpetgrass, dichondra, ground covers, garden plants, flowers and shrubs.

Do not apply to centipedegrass turf in the Spring before March 15 (or before the danger of frost has passed), or after September. Do not apply to bahia before March 15 or after November 1.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a clean, dry place. Reseal opened bags by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

**Suggested Spreader Settings**

to provide proper distribution calibrate spreader before application

61 7/8 lbs (28.07 kg) treats 22,000 ft<sup>2</sup> (1/2 acre/2044m<sup>2</sup>) at RECOMMENDED RATE.

61 7/8 lbs (28.07 kg) treats 22,000 ft<sup>2</sup> (1/2 acre/2044m<sup>2</sup>) at HALF RATE and reduced width. \*

<u>Spreader</u>	<u>Ground PTO Speed</u>	<u>Width of Coverage</u>	<u>Spreader Settings</u>	
			<u>Half*</u>	<u>Recommended</u>
Scotts (Drop type)	3 mph	overlap wheels	-	4 1/2
Scotts Rotaries	3 mph	7 feet *	B 1/2*	D 1/2
Lely TR & WFR models	4.5 mph	___ feet *	___ *	___
Lely - HR (PTO model)	450 rpm	___ feet *	___ *	___
Gandy (30")	3 mph	overlap wheels	-	24

\*At the HALF-RATE setting, apply at coverage width of 3-1/2 feet for Scotts rotaries and \_\_\_ feet for the Lely rotaries to achieve the overall RECOMMENDED RATE.

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (WH-567)  
WASHINGTON, D.C. 20460

FPA REGISTRATION NO.

DATE OF ISSUANCE

MAY 24 1984

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

NOTICE OF PESTICIDE:  REGISTRATION  
 REREГИSTRATION

(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (include ZIP code)

[ ] [ ]  
[ ] [ ]

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL


MAY 24 1984

ProTurf Fertilizer Plus Dicot Weed Control (XXXX)

7/18/83 ADR  
7/25/83 ADR  
7/27/83 ADR  
10/19/83 RF

FRONT:

(Scotts (R))  
ProTurf  
Brand



32-4-4

Fertilizer Plus  
Dicot Weed Control II

- . provides post emergence dicot weed control
- . provides controlled release nitrogen feeding

Net Weight 61 7/8 lbs (28.07 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs.

Environmental Hazards: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on the label.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid . . . . . 1.22%  
 2-(2-Methyl-4-chlorophenoxy) propionic acid . 1.22%  
 Dicamba: 2-Methoxy-3,6-dichlorobenzoic acid . . . . . 0.081%

INERT INGREDIENTS: . . . . . 97.479%

Total 100.00%