

PM

23

538-186

4/30/98

page 1 of 5

12/3/92 PG
6/17/93 VScr
3/10/98 VS/kjl

RANGE LABEL


FRONT:

ProTurf Fertilizer Plus Dicot Weed Control III

EPA Reg No 538-186 Stock #8609 (S2652)

(Scotts)
ProTurf
BRAND

XX-X-X
Fertilizer Plus
Dicot Weed
Control III



- provides postemergence dicot weed control
- provides controlled release nitrogen feeding

Net Weight 38.04 — 61.88 lb (17.24 — 28.06 kg)

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS**

Hazards to Humans and Domestic Animals: CAUTION, avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs.

Personal Protective Equipment

*waterproof is O.K. (WPS Guidance For 2,4-D)
chemical resistant category (gloves)*

Applicators and other handlers must wear protective eyewear, waterproof gloves, long-sleeved shirt, long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handlers PPE requirements may be reduced or modified in the WPS.

2 9 5

Environmental Hazards: Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on the label. Drift or runoff may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D/MCPP containing pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid.....	0.75 -	1.220%
2-(2-Methyl-4-chlorophenoxy) propionic acid.....	0.75 -	1.220%
Dicamba: 3,6-dichloro-o-anisic acid.....	0.050 -	0.081%
INERT INGREDIENTS:	98.450 -	97.479%
	Total	100.000%

EPA Reg No 538-186

EPA Est 538-OH-1

ProTurf Fertilizer Plus Dicot Weed Control III XX-X-X

643

Guaranteed Analysis

Total nitrogen (N).....	XX%
XX% ammoniacal nitrogen	
XX% urea, methylene ureas nitrogen	
XX% water insoluble nitrogen	
Available phosphate (P ₂ O ₅).....	X%
Soluble potash (K ₂ O).....	X%

Derived from monoammonium phosphate, urea, methylene ureas and potassium sulfate.

Product of USA

Professional Business Group
The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041

©1998. The Scotts Company. World rights reserved.

Recommended For Use By Professional Turfgrass Managers

8609 XX-X-X
Fertilizer Plus
Dicot Weed
Control III

GUSSETS:

XXXX	(Scotts®) ProTurf® BRAND	XX-X-X Fertilizer Plus Dicot Weed Control III
------	--------------------------------	--

538-186 Dicot III

BACK

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, ~~waterproof~~ gloves, shoes plus socks, protective eyewear.

waterproof gloves is O.K.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

ORNAMENTAL TURFGRASS USE

This product applies 1.5 pounds of 2,4-D per acre per application. The maximum number of broadcast applications per treatment site is two per year.

Not for use on golf course putting greens. Fertilizes grass and controls these rosette and vining type weeds:

- aster
- bittercress
- black medic
- bristly oxtongue
- buckhorn
- bull thistle
- burclover
- buttonweed
- carpetweed
- catsear (coast dandelion)
- chickweed
- chicory
- cinquefoil
- clover (white)
- cudweed
- curly dock
- dandelion
- dichondra

- English daisy
- eveningprimrose
- filaree
- fleabane
- ground ivy
- hawkweed
- healall
- henbit
- horseweed
- knotweed (seedling)
- lambsquarier
- lippia (matchweed)
- mallow
- moneywort
- morningglory
- mustard
- pennycort
- pepperweed

- pigweed
- plantain
- prickly lettuce
- purslane
- pusstoes
- ragweed
- scarlet pimpernel
- shadflower (whitlowort)
- shepherdspurse
- smartweed
- sowthistle
- stichwort
- threeseeded mercury
- thymeleaf sandwort
- vech
- yellow rocket
- yellow toadflax

For use on ornamental Turfgrass areas other than putting greens.

Apply to moist foliage at the RECOMMENDED RATE or for better distribution when using a rotary spreader, apply at the HALF RATE with a reduced width of coverage (see spreader settings). Apply when weeds are growing actively. Grass should not be so tall it touches the bottom of the spreader.

Rain, sprinkling, mowing, or heavy traffic within 24 hours after application can knock granules off weeds and may reduce effectiveness. Apply under non-windy conditions, thereby reducing drift to non-target areas, and optimizing particle distribution for good weed control. Retreating may be necessary, but wait at least 3 weeks to observe full results of first application.

Precautions

Avoid application when daytime temperatures are consistently 90°F or above.

Do not over apply, especially under the branch spread of trees and shrubs.

On new grass plantings delay application until after the fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 1 month after application.

Not for use on creeping bentgrass, St. Augustinegrass, carpetgrass, dichondra, ground covers, garden plants, flowers and shrubs.

Do not apply to centipedegrass turf in the Spring before March 15 (or before the danger of frost has passed), or after September. Do not apply to bahiagrass before March 15 or after November 1.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Professional Business Group
The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

This bag treats 22,000 ft² (1/2 acre) at RECOMMENDED RATE.
This bag treats 22,000 ft² (1/2 acre) at HALF RATE and HALF WIDTH.**

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE</u>	<u>HALF RATE/ HALF WIDTH**</u>
Scotts Drop Type	3 mph	overlap wheels	X	X
Gandy - 30"	3 mph	overlap wheels	X	X
Scotts Rotaries*				
R-7, R-7X	3 mph	XX feet	X	X
R-8, R-8A (cone 8)	3 mph	XX feet	X	X
Lely Models (see below)	4.5 mph	XX feet	X	X

* Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

** At the HALF RATE setting, apply at coverage width of 3 1/2 feet for Scotts rotaries and 9 feet for the Lely rotaries to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric coverage: 28.1 kg treats 2044m² at the RECOMMENDED RATE.

NOTE TO EPA: This product applies 1.5 lbs of 2,4-D, 1.5 lbs of MCPP and 0.1 lbs of Dicamba/acre per application at normal rate.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.

Professional Business Group
The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041