

FRONT:

185
1984 PACKAGE COPY FOR
LAWN INSECT CONTROL 40006

03/16/83 KGB 08/22/83 ER
03/30/83 KGB 11/21/83 AR
04/20/83 KGB
06/21/83 KGB

(Scotts (R)) Lawn Insect
Control

(NOTE: Above description will be inverted.)


Treats 5,000 sq ft (465 m²)

(Scotts (R))

Lawn Insect
Control

. WHITE GRUBS . CHINCH BUGS . SOD WEBWORMS

and other insects listed on the back.



NET WEIGHT 9 1/4 LBS (4.20 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION: Harmful if swallowed. May be absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid inhalation. If on skin: remove contaminated clothing and wash affected areas with soap and water. If in eyes: flush eyes with plenty of water. Call a physician if irritation persists. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs. Water treated area after application and allow to dry before permitting use.

Note to Physician: Diazinon is a cholinesterase inhibitor. Atropine is antidotal if symptoms of cholinesterase inhibition are present.

BEST DOCUMENT AVAILABLE

537-117

26

FRONT CONTINUED:

Environmental Hazards: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bags by folding top down and securing.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

ACTIVE INGREDIENT:

Diazinon [O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate] 5.84%

INERT INGREDIENTS: 94.16%

Total 100.00%

Product of USA

EPA Reg No 538-XX LPA Est 538-OH-1

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1984, The O M Scott & Sons Company. All rights reserved.

40006 (Scotts (R)) Lawn Insect Control

BESLI

52-107

(2)

RIGHT GUSSET:

(Scotts(R)) Lawn Insect
Control

40006

LEFT GUSSET:

40006

(Scotts (R)) Lawn Insect
Control

337-111

BACK:

40006 LIC 5M
EASY
OPEN BAG
PULL
WHITE
STRING

Treats 5,000 sq ft (465 m²)

(Scotts(R))

Lawn Insect
Control

Billbugs

Sod webworms

Mole crickets

Chinch bugs

White grubs

Also controls the following insects:

- | | | |
|-----------------|-----------------|-----------------------|
| . armyworms | . cutworms | . brown dog ticks |
| . Bermuda mites | . earwigs | . sod webworms |
| . billbugs | . fleas | (lawn moth larvae) |
| . chiggers | . flea beetles | . vegetable weevil |
| . chinch bugs | (in dichondra) | (in dichondra) |
| . clover mites | . leafhoppers | . white grubs such as |
| . crickets | . mole crickets | Japanese beetle, |
| | | European chafer, |
| | | Masked chafer, |
| | | Phyllophaga (May or |
| | | June beetle) |

Easy to apply

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- Apply when insect activity or damage (chewed or brown grass blades) is first observed in the lawn.
- Apply when lawn is dry.
- Water lawn after application to wash granules from the lawn foliage. If treating for grubs, water thoroughly (so active ingredients will be carried into the root zone.). Don't allow traffic on the lawn until it has dried.
- Treat the entire lawn to prevent further insect damage.
- You should not use any other lawn control product for at least one week after application.
- A second application may be needed in 6 weeks if insect activity continues.
- Do not use on fruit or vegetable plants.

BEST DOCUMENT AVAILABLE

BACK CONTINUED:

GRUB CONTROL - The most likely period for grubs is late July through early October, but in some cases grubs are active in the Spring. An application for Spring grub activity requires a repeat Summer - early fall application for effective control.

For best results - Water your lawn 24 hours prior to application. This will better enable the active ingredients to penetrate the thatch and control grubs.

CHINCH BUGS - Start treatments when daily high temperatures are 80° or above. A second application may be needed in 6 weeks if insect activity continues.

MOLE CRICKETS - The best time to apply is in May to July when the insects are young.

For best results - Water your lawn 24 hours prior to application. This draws insects closer to the surface for better control.

Apply at Scotts Spreader (PF, Standard or Automatic)
setting X
(treats 5,000 sq ft)

TIPS ON SPREADING

For best results we recommend Scotts drop type spreaders.

- First apply 2 header strips (ILLUS.)
- Operate spreader the longest way of the lawn
- Overlap wheels to insure complete insect control (ILLUS.)
- Shut off spreader when you have reached the header strips or anytime you stop

40006

Scotts Policy
No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTTS...the lawn people
MARYSVILLE, OHIO 43041

0 (UPC)...

32247 40006

40006

(Scotts (R))

Lawn Insect
Control

00380