

Mr. Vincent Snyder, Jr.
O.M. Scott and Sons Company
14310 Scottlawn Road
Marysville, OH 43041

FEB 15 1989

3/10 / 17 : 219898 / 1

Dear Mr. Snyder:

Subject: Amendment - Confidential Statement of Formula
With Alternate Carrier and Source of Inert
Ingredient and Label Revision
Lawn Insect Control
EPA Registration No. 538-187
Your Applications Dated March 29 and June 20, 1988

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable provided that you make the labeling change indicated below prior to release for shipment of product bearing the amended labeling.

- Update the Environmental Hazards statement by changing "cleaning of equipment or disposal of wastes" to "disposing of equipment washwaters."

The alternate Confidential Statement of Formula is acceptable and has been made a part of the record for this product.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure
50711:1:Richards:LLR-13:KENCO:2/14/89:3/29/89:rw:ek:rv

CONCURRENCES								
SYMBOL								
SURNAME								
DATE								

FRONT:

FOR EPA APPROVAL
1988 PACKAGE COPY FOR
LAWN INSECT CONTROL 40006
02/18/88 AR 3/2/88 AR

(Scotts) Lawn Insect
Control

(NOTE: Above description will be inverted.)

Treats 5,000 sq ft (465 m²)

SCOTTS POLICY
No-Quibble
Guarantee
SEE BACK FOR DETAILS

(Scotts(R))

Lawn Insect
Control

controls white grubs
chinch bugs
sod webworms
European crane fly larvae
and other insects listed on the back.

This product must not be used on golf courses and sod farms

NET WEIGHT 9 1/4 LBS (4.20 kg)

LEFT COLUMN

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Under the Federal Insecticide, Fungicide, and Rodenticide Act
as amended for the purpose
of...
526-187

Hazards to Humans and Domestic Animals: CAUTION: Harmful if swallowed. May be absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid inhalation. If on skin: remove contaminated clothing and wash affected areas with soap and water. If in eyes: flush eyes with plenty of water. Call a physician if irritation persists. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs. Water treated area after application and allow to dry before permitting use.

Note to Physician: Diazinon is a cholinesterase inhibitor. Atropine is antidotal if symptoms of cholinesterase inhibition are present.

3/7

FRONT CONTINUED:

Environmental Hazards: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes,

RIGHT COLUMN

streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENT:

Diazinon [0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate]

5.84%

INERT INGREDIENTS: 94.16%

Total 100.00%

EPA Reg No 538-187

EPA Est 538-OH-1

Product of USA

The O M Scott & Sons Company
Marysville, Ohio 43041

(c) 1988, The O M Scott & Sons Company. All rights reserved.

(Printing Calendar)

GUSSETS:

4
7

RIGHT GUSSET:

40006 (Scotts) Lawn Insect
Control

SCOTTS POLICY
No-Quibble
Guarantee
SEE BACK FOR DETAILS

LEFT GUSSET:

(Scotts) Lawn Insect
Control

SCOTTS POLICY
No-Quibble
Guarantee
SEE BACK FOR DETAILS

40006

BACK:

PULL TAB
and remove for
pour spout

(ILLUS.)

Easy
Open PULL HERE

Treats 5,000 sq ft (465 m²)

LIC - 40006
5M
PROOF
OF
PURCHASE
(Scotts)

(Scotts(R))

Lawn Insect
Control

Controls the following insects:

ants	cutworms	brown dog ticks ;
armyworms	earwigs	sod webworms
Bermuda mites	European crane fly larvae	(lawn moth larvae)
billbugs	(leatherjackets)	vegetable weevil
chiggers	fire ants (except in	(in dichondra)
chinch bugs	California)	white grubs such as
clover mites	fleas	Japanese beetle,
crickets	flea beetles	European chafer,
	(in dichondra)	Masked chafer,
	leafhoppers	Phyllophaga (May or
	mole crickets	June beetle)

Easy to Apply

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- Apply when insect activity or damage (chewed or brown grass blades) is first observed in the lawn.
- Apply when lawn is dry.
- Water lawn after application to wash granules from the lawn foliage. If treating for grubs and mole crickets, water thoroughly (so active ingredients will be carried into the root zone). Don't allow traffic on the lawn until it has dried.
- Treat the entire lawn to prevent further insect damage.
- You should not use any other lawn control product for at least one week after application.
- A second application may be needed in 6 weeks if insect activity continues.
- Do not use on fruit or vegetable plants.

BACK CONTINUED:

6
7

GRUBS--

Northern States--The most likely period for grubs is late July through September but in some cases grubs are active in the spring. An application for spring grub activity requires a repeat summer-early fall application for effective control.

Southern States--Apply during June-August. A second application may be needed six weeks later. Applications made later in the year, when grubs are more mature, are less effective.

For best results - Water your lawn 24 hours prior to application. This will better enable the active ingredients to penetrate the thatch and control grubs.

CHINCH BUGS-- Start treatments when daily high temperatures are 80° or above. A second application may be needed in 6 weeks if insect activity continues.

MOLE CRICKETS-- The best time to apply is in May to July when the insects are young.

For best results - Water your lawn 24 hours prior to application. This draws insects closer to the surface for better control.

CRANE FLY LARVAE (leatherjackets)--Apply when activity is first noticed in the spring, usually mid-March to mid-April.

FIRE ANTS (except in California) - Apply 1/2 cup of product on each fire ant mound, water gently and thoroughly after application. For best results apply in early morning or late evening.

Application Rate	Scotts Drop Spreaders	Scotts EasyGreen™ Rotary
(treats 5,000 sq ft)	5	25 1/2

TIPS ON SPREADING WITH SCOTTS SPREADERS

Scotts Drop:

- First apply 2 header strips (ILLUS.)
- Operate spreader the longest way of the lawn
- Overlap wheels to insure complete insect control
- Shut off spreader when you have reached the header strips or anytime you stop (ILLUS.)
- Mow lawn first if grass is tall enough to touch the bottom of spreader

EasyGreen™ Rotary:

- Slide shut-off lever to the right. Application automatically starts and stops when you do
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs

BACK CONTINUED:

7
-7

40006

**Scotts Policy
No-Quibble Guarantee**

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

(Scotts®) the lawn people
Marysville, Ohio 43041

0 (UPC) 0

32247 40006

(NOTE: The description below will be inverted.)

40006

(Scotts) Lawn Insect
Control

0038o