



# (Scotts.)

## Lawn Insect Control Control





- Kills-lawn-damaging insects that cause your lawn to thin and turn brown
- · Kills fleas and Brown Dugticks before they get on your per

#### KEEP OUT OF REACH OF CHILDREN CAUTION

See back for additional precautionary statements.

ACTIVE INGREDIENT: Distinont . Total 100 00% EPA Reg. No. 538-187 EPA Est. No. 538-0H-1 (S) EPA Est. No. 44616-MO-01 (H) ATCASS/833-41-5 # Subscript used is first letter of run code on end of container Product of U.S.A. Treats 5,000 sq. ft.

NET WEIGHT 10.10 b (4.58 kg

### Controls these insects

ants armyworms Bermudagrass mites billbugs chiggers chinch bugs

clover mites

crickets cutworms earwigs


European crane fly larvae (leatherjackets) fire ants

(except in California)

flea beetles (in dichondra) leafhoppers mole crickets Brown dog ticks sod webworms

(lawn moth larvae)

vegetable weevil (in dichondra) white grubs such as Japanese beetle, European chafer, Masked chafer Phyllophaga (May or June beetle)



#### KEY PRODUCT FACTS

#### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling

  Apply product to entire lawn when insect activity or damage (chewed or browned)
- leaves) is first observed in the lawn.

  Apply product to a dry lawn.

  Water-in product after application. Don't allow traffic on the lawn until it has dried.
- Do not use any other weed or disease control products for at least one wee after application.
- For severe insect problems, a second application may be needed in 6 weeks.
   Do not use on fruit or vegetable plants.
   FOR FIRE ANTS (except in California).

- Apply 1/2 cup of product on each fire ant mound.

  Water gently and thoroughly after application.

  For best results, apply in early morning or late evening. • FOR CHINCH BUGS
- Start treatments when daily high temperatures are 80°F or above and repeat at 6 week intervals if insect activity continues.
  FOR MOLE CRICKETS
  - Water your lawn first, let lawn dry, and then apply product. Water in product thoroughly. Apply in May to July, when insects
- are young.
   FOR CRANE FLY LARVAE (leatherjackets)
  Apply when activity is first noticed in the spring, usually mid-March to mid-April.



#### Special Precautions to Reduce Risks to Birds

Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

- a) Do not treat areas larger than 15,000 sq. ft.
  b) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
- c) Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
- d) Avoid overlapping treated areas. e) Follow all watering instructions included within the directions for use, however, stop watering before puddling occurs.
- f) Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
- g) If necessary to repeat application, wait a minimum of 3 weeks before applications. Make no more than 4 applications per year.

#### Socrading Tips apply r con longest (g) of lawπ.

#### Spreader Settings

For hest completely seems production in a grant before applying product.

#### Scottlerentry Spreaders

Secret S Speedy Green® Make cockeres chant 5 feet from previous pass to overlap

Scotts'3 EasyGreen®

rate elen passanta sometes. us pass to overlap





kopute oprender, wheel completely overlaps previous

Scotts\*\*\*4 PrecisionGreen®

Completely overlaps previous 63/4

Additional Spreader Settings:

EZ Drop, Republic (#530) ..... 

Scotts Broadcast Tractor Spreader .....

(Non-Scotts spreader settings are approximate You may need to adjust for proper coverage.)

#### First Aid and Environmental Information

#### KEEP OUT OF REACH OF CHILDREN CAUTION

#### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin, or clothing. Wash thoroughly after breathing dust and contact with eyes, skill, or crouling, wash stondager, handling and before smoking or eating. Do not store near food or feed. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until grapules have been watered in and the grass or soil has dried

First Aid

If swallowed: Catfa physician or Poison Control Center immediately. Drink one or two glasser of writer and induce vomiting by touching the back of the throat with finger. Rej cat uctil vomit fluid is clear. Do not induce vomiting or give

anything by mouta't person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. If on skin: Remove contaminated clothing and avash affected areas with soap and water.

If in eyes: Rinse eyes vising entry of water and call a physician immediately. Note to Physician: This product condite at the granophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be a ministered, but only in conjunction with atropine.

40006-7

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of granules. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. contaminate water by cleaning of equipment or disposal of equipment washwaters. Storage and Disposal

STORAGE: Store in original container in a cool, dry, well-ventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Reseal opened bag by folding top down

and securing PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash

CONTAINER DISPOSAL: Do not reuse empty bag, Discard bag in trash. Distributed by: The Scotts Company, 14111 Scottslawn Rd., Marysville, OH 43041

#### Scotts No-Quibble Guarantee

If for any reason you kit. Lok satisfied with results after using this product, you are crititled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

Product of U.S.A. ຂ 2000, The Scotts Contoling

Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-sc.com">http://www.regulatory-info-sc.com</a>.

#### Questions / Information



Consumer Helpline 1-800-543-TURF (8873)



Visit our website www.scottslawn.com



Scotts Company 14111 Scottslawn Rd. Marysville, Ohio 43041

