

538-188

PM25

1 of 5

FRONT:

PROTURF SOUTHERN WEED-
GRASS CONTROL #83204
08/15/84 RF 01/04/85 MJB
09/19/84 RF 05/29/85 JF
10/09/84 RF 06/24/85 JF

(SCOTTS(R))
ProTurf(R)
BRAND

Southern
Weedgrass
Control

- preemergent control of goosegrass, crabgrass, Poa annua and other annual grassy weeds in southern grasses
- preemergent control of several broadleaf weeds

Net Weight 28 1/4 lbs (12.81 kg)

LEFT COLUMN

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

ACCEPTED

OCT 23 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 538-188

Hazards to Humans and Domestic Animals: CAUTION. Causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation.

Environmental Hazards: This pesticide is toxic to fish. Do not apply



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

GUSSETS:

RIGHT GUSSET

83204

(Scotts(R))
ProTurf(R)
BRAND


Southern
Weedgrass
Control

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

Southern
Weedgrass
Control

83204
(Packaging Calendar)



4065

BACK:

Southern
Weedgrass
Control

83204

(SCOTT'S (R))
ProTurf (R)
BRAND

Southern
Weedgrass
Control

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls these grassy weeds as they sprout:

- | | |
|------------|---------------|
| goosegrass | Poa annua |
| crabgrass | barnyardgrass |
| foxtail | fall panicum |

Controls these broadleaf (dicot) weeds as they sprout:

- | | |
|------------|-----------|
| chickweed | evening- |
| cudweed | primrose |
| hop clover | prostrate |
| henbit | spurge |
| oxalis | |

For use on established bermudagrass (non-winter overseeded), St. Augustinegrass, centipedegrass, zoysiagrass, bahiagrass, and tall fescue.


For goosegrass, crabgrass, and other annual grasses, and germinating dicot weeds, apply in the spring before they sprout. For Poa annua, chickweed, and henbit apply in late summer-early fall before weeds sprout.

Water after application (1/4 to 1/2 inch) to remove granules from the foliage and to move the product to the soil surface. Do not destroy the chemical barrier after application by raking or aerifying.

Do not contaminate feed or foodstuffs. Keep children and pets off treated areas until material is washed into soil and foliage has dried.

Note Carefully

Not for use on cool season grasses or dichondra.



BACK CONTINUED:

RIGHT COLUMN

This product should not be used on winter overseedings.

This product should not be used on newly seeded or sprigged turfgrass areas or where turf has been severely thinned due to winter damage. Wait until the stand is uniform and dense.

Residual effectiveness requires a delay of four growing months after treatment before reseeding or winter overseeding and five growing months before sprigging.

Product does not control germinated or established weeds.

Limit use to twice yearly allowing at least four months between applications.

Storage and disposal:

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

28 1/4 lbs treats 11,000 ft² (1/4 acre) at RECOMMENDED RATE

<u>Spreader</u>	<u>Ground Speed</u>	<u>Width of Coverage</u>	<u>Recommended Rate</u>	<u>Half Rate/ Half Width*</u>
Scotts (Drop type)	3 mph	overlap wheels	6 1/4	not recommended
Gandy (30")	3 mph	overlap wheels	31	not recommended
Scotts Rotaries				
R-7, R-7X	3 mph	8 feet	F 1/2	D 1/2 1/4 slide
R-8 (cone)				
Lely-(Models, see below)	4.5 mph	22 feet	4 3/4 III	4-111

* At the Half Rate/Half Width setting apply at a swath width of 4 ft for the Scott Rotary and 11 ft for the Lely to achieve the overall Recommended Rate.

Metric application rate: 12.81 kg treats 1,022 m² at the RECOMMENDED RATE

Lely Models WTR, WRF, HR, and 1250. PTO at 450 rpm

For proper distribution calibrate your spreader before application