

PM 25 538-190

1042



Treats up to 5,000 sq. ft.

CRABGRASS PREVENTER PLUS FERTILIZER



Treats up to 5,000 sq. ft.
CRABGRASS PREVENTER PLUS FERTILIZER

ACCEPTED
APR - 5 1995
United States Federal Insecticide, Fungicide, and Rodenticide Act
as amended, for the pesticide registered under
EPA Reg. No. **538-190**

21-3-3

FEATURING

Scotts

Slow-release nitrogen fertilizer - for uniform growth and extended feeding



Scotts



41005

NET WEIGHT
19.0 LBS (8.61 kg)

PREVENTS CRABGRASS



ACTIVE INGREDIENTS:
Fertilizer 21-3-3
2,4-D (2,4-Dichlorophenoxyacetic acid) 0.01%
NITROGEN 21.00%
PHOSPHORUS 3.00%
POTASSIUM 3.00%
EPA Reg. No. 538-190 EPA Est. No. 6379-B-1
*Licensed under one of the terms of the following U.S. Patent: 3,629,722-4,581,241
CAUTION: KEEP OUT OF REACH OF CHILDREN.
See back for additional precautionary statements.

BEST COPY AVAILABLE

1042

U 32247 41005 2

Treats up to 5,000 sq. ft.
**CRABGRASS PREVENTER
PLUS FERTILIZER**



Scott's

41005

TECHNOLOGY

POSSIBLE PREVENTION

FEATURING

CRABGRASS PREVENTER PLUS FERTILIZER

21 • 3 • 3

CRABGRASS PREVENTER PLUS FERTILIZER 21 • 3 • 3
Guaranteed Analysis

Total Nitrogen (N)	21%
0.8% water-soluble nitrogen	
20.2% urea nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Phosphoric Acid (P ₂ O ₅)	3%
Sulfur (S) (lb/100)	2.5%
Derived from: diammonium phosphate, potassium chloride, polymer-encapsulated sulfur-coated urea and urea.	
Contains 5.0% slowly available feeding nitrogen from polymer-encapsulated sulfur-coated urea.	
U.S. Patent No. 5,219,465	
Product of U.S.A.	

Manufactured by The Scotts Company, 10111 Southway Rd., Maryville, TN 37804

CAUTION: KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS: CAUTION. Causes eye irritation. Do not get in eyes, on skin, or clothing. If in eyes or on skin, flush with plenty of water and get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Do not consume food or beverages. Do not drink treated water. Do not feed animals to livestock.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Avoid areas where water may be present in treated areas. Do not apply to waterways or streams. Do not apply to areas where water may be present in treated areas. Do not apply to areas where water may be present in treated areas.

STORAGE: Store in a clean, dry place. Protect against fire by keeping away from heat and flames.

PESTICIDE DISPOSAL: See label for disposal instructions in several states of use and disposal in trash.

CONTAINER DISPOSAL: Do not reuse bag. Recycle bag in trash.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

APPLICATION

Apply when your lawn is dry. If it is necessary to water this product in after application.

Apply in early spring before the third or fourth mowing, or before summer rains are regular in the fall, late spring, or before winter rains will affect the performance of this product.

- For Kentucky bluegrass, fine fescue, tall fescue, perennial ryegrass, and blends of these grasses, apply at the RATE.
- For bermudagrass, St. Augustinegrass, zoysiagrass, centipedegrass, and blends of these grasses, apply at the RATE.

Apply to prevent germination of crabgrass, foxtail, barnyardgrass, tall fescue, goosegrass, and other weeds (not for use on annual ryegrass).

- In early spring before the third or fourth mowing, or before summer rains are regular in the fall, late spring, or before winter rains will affect the performance of this product.
- For heavy infestations, a repeat treatment is recommended after 6 to 8 weeks.

Apply in the fall (August-October), after temperatures have cooled and but before heavy frosts, to prevent weeds from becoming established during winter or next spring, or to prevent germination of fall annuals and biennials.

USE PRECAUTIONS
Do not seed or soil for 4 weeks following application. The pre-emergent barrier prevents grass seed from germinating.

Not for use on driveways or hard-paved areas.

Except for heavy infestations of weeds, allow at least a 4-month interval between applications of this product. Do not apply this product more than once annually.

WARNING: Do not walk or drive for a month after application. To avoid disturbing the pre-emergent barrier on the soil surface.

*Holding water: Can cause staining to concrete, stone, asphalt surfaces (such as sidewalks and garden footings, steps, or stairs). If water is spilled, brush off particles to prevent staining and/or staining into the floor. If necessary, rinse off with water.

SPREADER SETTINGS

For best results, use a Scott's spreader.

	Regular Rate (lb/1000 sq. ft.)	Heavy Rate (lb/1000 sq. ft.)
Scott's Drop Spreader	6	8 1/2
Scott's EasyGreen® Rotary	20	25 1/2
Scott's SpeedyGreen® Rotary	4	4 1/2

Additional Spreader Settings

EZ Drop Spreader (E330) or Precision	8 1/2
EZ Broadcast Spreader (E330) or Precision	2 1/2
Rot Dye Drop (E330-72)	11
Rot Dye Rotary (E330-50)	1
Quaker Drop Spreader (E330-50)	6
Quaker Rotary Spreader (E330-50)	4 1/2

Note: Scott's spreader settings are approximate. You may need to adjust for proper coverage.

Scott's Drops

- First apply 2 heaping drops.
- Operate spreader the longest way of the wheel.
- Overlap wheels to insure uniform feeding.
- Stop all spreader when you have reached the heaping drops or stopped.

EasyGreen® Rotary

- Slide shut-off lever to the right, position automatically starts and runs when you go.
- Adjust wheel by turning each pair about 1/8 inch from the previous one to insure overlapping areas.

SpeedyGreen® Rotary

- Slide shut-off lever to the right, position automatically starts and runs when you go.
- Adjust wheel by turning each pair about 1/8 inch from the previous one to insure overlapping areas.

©1991, The Scotts Company. Scott's is a registered trademark.

QUESTIONS?

Call Scott's toll-free Customer Hotline
1-800-543-TURF

For additional information, please
visit our website at www.scotts.com